INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps.

Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6" x 9" black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.

Bell & Howell Information and Learning
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA

UMI[®]
800-521-0600

Dynamics of Harpooning Studied by
Transition State Spectroscopy

by

Andrew J. Hudson

A thesis submitted in conformity with the requirements
for the degree of Doctor of Philosophy,
Graduate Department of Chemistry, in the
University of Toronto.

© Copyright by Andrew J. Hudson

National Library
of Canada

Acquisitions and
Bibliographic Services

395 Wellington Street
Ottawa ON K1A 0N4
Canada

Bibliothèque nationale
du Canada

Acquisitions et
services bibliographiques

395, rue Wellington
Ottawa ON K1A 0N4
Canada

Your file *Votre référence*

Our file *Notre référence*

The author has granted a non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of this thesis in microform, paper or electronic formats.

The author retains ownership of the copyright in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de cette thèse sous la forme de microfiche/film, de reproduction sur papier ou sur format électronique.

L'auteur conserve la propriété du droit d'auteur qui protège cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

0-612-41440-X

Thesis Abstract

Dynamics of Harpooning Studied by Transition State Spectroscopy

by

Andrew J. Hudson

Department of Chemistry, University of Toronto

1998

The van der Waals complexes, $\text{Na}\cdots\text{XR}$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}, \text{R}=\text{CH}_3$; and $\text{X}=\text{F}, \text{R}=\text{H}$), have been generated by crossing a beam of sodium with the expansion region of a supersonic jet of the appropriate halide in an inert-carrier gas. The complexes were identified by photoionization time-of-flight mass spectrometry. Depletion of the complexes was observed following photoexcitation using a broad range of wavelengths, λ_1 . Initial excitation of the Na chromophore within the complex is believed to be followed by charge-transfer dissociation;

An experimental study of the photodepletion of $\text{Na}\cdots\text{XR}$ complexes has been made, and the results have been compared to recent *ab initio* studies of the same process. Photoexcitation of $\text{Na}\cdots\text{XR}$ accesses a potential-energy surface (PES) for the reaction $\text{Na}^*+\text{XR}\rightarrow\text{NaX}+\text{R}$, at a configuration intermediate between the reactants, Na^*+XR , and products, $\text{NaX}+\text{R}$. We, therefore, denote the excited-state species as a transition state, TS, indicated by the $\ddagger$ symbol. The starting configuration for the TS, $[\text{Na}^*\cdots\text{XR}]^\ddagger$, is determined by the geometry of $\text{Na}\cdots\text{XR}$ and the excitation energy, $h\nu_1$. Consequently, by varying the wavelength, λ_1 , we are able to access selected TS configurations. This constitutes a Transition State Spectroscopy for these classic 'harpooning' reactions. As the excitation energy is further increased, we are able to access the TS for the reaction of XR with successively higher electronically-excited states of Na^* .

The dynamics of the excited-state reaction have been probed by measuring the depletion cross-section for the ground-state complex as a function of the excitation wavelength. In addition, the lifetime of the TS was estimated by measuring an ion signal for photoionization directly from the TS.

The TS is believed to dissociate via surface-hopping from the electronically-excited state into alternate channels on the ground-state PES. The surface hop occurs in the region in which the character of the TS changes from largely covalent to largely ionic, $[\text{Na}^* \cdots \text{XR}]^\ddagger \rightarrow [\text{Na}^+ \cdots \text{XR}^-]^\ddagger$; this is the harpooning event proposed over half a century ago to be the crucial step in the reaction of alkali metals with halogens.

Acknowledgments

I sincerely thank John Polanyi for his support during the work described in this thesis. It has been a memorable experience working under his guidance.

I am greatly indebted to Jixing Wang and Rudolf Ehlich for teaching me experimental techniques and for their participation in the measurements of the $\text{Na}\cdots\text{XCH}_3$ ($\text{X} = \text{F}, \text{Cl}, \text{Br}$) and $\text{Na}\cdots\text{FH}$ photodepletion spectra. I would like to thank Han Bin Oh for his participation in the measurement of the Transition State ionization spectrum for $[\text{Na}^*\cdots\text{FH}]^\ddagger$.

This work was made especially interesting as a result of a successful collaboration with theoreticians from our group. Xiao Yan Chang provided *ab initio* calculations of the minimum-energy structures, binding energies and vibrational frequencies for $\text{Na}\cdots\text{XCH}_3$ and $\text{Na}^*\cdots\text{XCH}_3$ complexes. Piotr Piecuch provided detailed *ab initio* calculations of the complete potential-energy surface for the ground and electronically-excited states of NaFH .

An analytical fit to the *ab initio* potential energies for NaFH calculated in this laboratory was obtained by Don Truhlar and Maria Topaler at the University of Minnesota. From this analytical fit they were able to calculate the vibrational-energy levels and Franck-Condon factors for vertical transitions and, thereby, simulate a photoabsorption spectrum for $\text{Na}\cdots\text{FH}$. Their results contributed to a greater understanding of the experimental spectra.

I would like to thank Jamie Donaldson for many helpful discussions and in particular for leading us to identifying a Transition State lifetime for $[\text{Na}^*\cdots\text{FH}]^\ddagger$ from measurements of the depletion intensity as a function of the excitation-laser fluence. Also, thanks to Giacinto Scoles for noticing that the Na beam was a supersonic expansion and correcting our value for the collision energy of the molecular beams.

Finally, I would like to acknowledge British Gas for a Research Scholarship and the University of Toronto for an International Student Award. The experiments were financed by the National Science and Research Council (NSERC) and Photonics Research Ontario (PRO).

Table of Contents

Chapter 1	Introduction	1
1.1	Photoinduced reaction in weakly bound systems	1
1.2	Transition State Spectroscopy	7
1.3	Harpooning reactions	9
	• <i>Reaction of Na atoms with methyl halide molecules</i>	11
	• <i>Reaction of Na atoms with HF molecules</i>	15
1.4	Summary and thesis preview	21
Chapter 2	Experiment	24
2.1	Experimental apparatus	26
	• <i>Vacuum system</i>	26
	• <i>Crossed-molecular beam assembly</i>	26
	• <i>Excitation laser, L_1</i>	33
	• <i>Ionization laser, L_2</i>	35
	• <i>Time-of-flight mass spectrometer</i>	38
	• <i>Timing circuitry</i>	45
2.2	Data Analysis	48
	• <i>Mass spectra</i>	48

• <i>Photodepletion measurements</i>	50
2.3 Summary	53
Chapter 3 Photodepletion of $\text{Na}\cdots\text{XCH}_3$; X=F, Cl, Br	54
3.1 Identification of complexes	55
3.2 Structure of complexes	64
3.3 Photodepletion measurements	67
3.4 Discussion	77
• <i>peak assignment</i>	77
• <i>peak intensity</i>	87
• <i>vibrational structure</i>	93
3.5 Summary	101
Chapter 4 Photodepletion of $\text{Na}\cdots\text{FH}$	105
4.1 Identification of $\text{Na}\cdots\text{FH}$ complexes	106
4.2 Photodepletion measurements	109
4.3 Discussion	115
• <i>Dynamics of the photoinduced reaction</i>	115
• <i>Vibrational structure and intensity of the photodepletion peaks</i>	127
4.4 Summary	133

Chapter 5 Transition State ionization of Na$\cdot$-FCH₃	135
5.1 Experiment	137
5.2 Results	140
5.3 Discussion	143
5.4 Summary	147
Chapter 6 Conclusions and future directions	150
6.1 Conclusion	149
6.2 Future directions	152
Thesis Summary	158
References	162

List of figures

Figure 1.1:	The covalent, $M+X_2$, and ionic, $M^++X_2^-$, interaction potentials for the alkali metal and halogen molecule.	10
Figure 2.1:	Side view of the vacuum system.	25
Figure 2.2:	Crossed-molecular beam assembly.	28
Figure 2.3:	TOF mass spectrum obtained by expansion of a mixture of CH_3OH and H_2O in Ar-carrier gas through the pulsed valve.	32
Figure 2.4:	Top view of the reaction chamber indicating the path of the excitation laser, L_1 , and the ionization laser, L_2 .	34
Figure 2.5:	TOF mass spectrum obtained by expansion of a 2.5% mixture of HF in He at 250kPa backing pressure through the pulsed valve.	36
Figure 2.6:	Power dependence of the ionization signal for $Na\cdot FH$ measured at 248nm.	39
Figure 2.7:	Power dependence of the depletion signal for $Na\cdot FH$ measured at $\lambda_1 = 660$ nm.	40
Figure 2.8:	The ion-signal intensity for $Na\cdot FH$ as a function of the time delay between the ionization laser, L_2 and the onset of the pulsed-extraction field.	42
Figure 2.9:	The ion-signal intensity for $Na\cdot FH$ as a function of the backing pressure for the pulsed valve and the time delay between the ionization laser, L_2 , and the pulsed valve.	47

Figure 2.10:	Normalised ion-signal intensity for Na $\cdot$ FH as a function of the time delay between the ionization laser, L ₂ , and the excitation laser, L ₁ .	49
Figure 3.1:	TOF ion spectra for Na+XCH ₃ , X=F,Cl,Br,I.	58
Figure 3.2:	Power dependence of the ion signals for Na, NaI and Na ₂ I in the TOF spectra.	62
Figure 3.3:	Geometry and dissociation energy of the most stable configurations for Na $\cdot$ XCH ₃ , X=F, Cl, Br, I.	66
Figure 3.4:	Action spectra for Na $\cdot$ XCH ₃ , X=F, Cl, Br.	68
Figure 3.5:	Birge-Sponer plots for the vibrational progressions observed in the experimental action spectra.	75
Figure 3.6:	Calculated electron-density distribution for the HOMO in the ground (3s ² S) and the excited (3p ² P _x , 3p ² P _z and 4s ² S) states of Na $\cdot$ FCH ₃ .	78
Figure 3.7:	Qualitative potential-energy profiles referring to the Na $\cdot$ X bond distance for the three complexes.	86
Figure 3.8:	Geometry and dissociation energy of the most stable configurations for the excited states of Na $\cdot$ FCH ₃ .	94
Figure 3.9:	Geometry and dissociation energy (D) of the most stable configurations for Na $\cdot$ XCH ₃ , X=Cl, Br.	97
Figure 4.1:	The dependence of the complex-ion signal for Na $\cdot$ FH on the composition and identity of the carrier gas in the expansion mixture.	107
Figure 4.2:	Action spectrum for Na $\cdot$ FH.	110

Figure 4.3:	Experimental and calculated power dependence of the depletion signal for the complex Na $\cdots$ FH at an excitation wavelength, λ_1 , of 660nm.	114
Figure 4.4:	<i>Ab initio</i> MRDCI calculation of the geometry and dissociation energy of the most stable configurations of NaFH in the ground and excited states.	118
Figure 4.5:	Adiabatic collinear PESs of the ground and first-excited states of NaFH generated by MRDCI calculation.	121
Figure 4.6:	Potential-energy curves for the ground and the electronically-excited states, along an approximate minimum-energy path for the collinear PES.	123
Figure 4.7:	(a) Experimental photodepletion spectrum for Na $\cdots$ FH. (b) Calculated absorption spectrum for Na $\cdots$ FH at 250K. (c) Calculated components of the absorption cross section.	128
Figure 5.1:	Top view of the vacuum system.	139
Figure 5.2:	The photodepletion spectrum and TS ionization spectrum for the Na $\cdots$ FCH ₃ complex.	141
Figure 5.3:	The dependence of the complex-ion signal for Na $\cdots$ FCH ₃ on the lifetime of the TS, [Na $\cdots$ FCH ₃] ‡ .	146
Figure 6.1:	MRDCI calculation of the potential energy curves for the ground state and excited states of NaF.	154

List of tables

Table 2.1:	Electrode voltages for the TOFMS	44
Table 3.1:	The spacing between the first vibrational transitions observed in each progression.	73
Table 3.2:	The anharmonicity constant observed in each vibration progression.	76
Table 3.3:	Location of peaks in the action spectrum and the calculated vertical transition energies to lowest excited states for Na $\cdot$ FCH ₃ .	80
Table 3.4:	Location of peaks in the action spectrum and the calculated vertical transition energies to lowest excited states for Na $\cdot$ ClCH ₃ .	83
Table 3.5:	Location of peaks in the action spectrum and the calculated vertical transition energies to lowest excited states for Na $\cdot$ BrCH ₃ .	84
Table 3.6:	Integrated depletion cross-section from the Na $\cdot$ FCH ₃ action spectrum and the calculated oscillator strength for each electronic transition.	89
Table 3.7:	Integrated depletion cross-section from the Na $\cdot$ ClCH ₃ action spectrum and the calculated oscillator strength for each electronic transition.	91
Table 3.8:	Integrated depletion cross-section from the Na $\cdot$ BrCH ₃ action spectrum and the calculated oscillator strength for each electronic transition.	92
Table 3.9:	Calculated frequencies for vibrational modes in ground and excited states of Na $\cdot$ FCH ₃ .	95

Table 3.10: Calculated frequencies for vibrational modes in ground and excited states of Na··ClCH ₃ .	98
Table 3.11: Calculated frequencies for vibrational modes in ground and excited states of Na··BrCH ₃ .	99
Table 4.1: Calculated frequencies for vibrational modes in ground and excited states of Na··FH.	132

Chapter 1

Introduction

1.1 Photoinduced reaction in weakly bound systems

In recent years there has been considerable interest in the study of photoinduced reactions in molecules ordered at surfaces[1] or in gaseous complexes[2]. In gaseous complexes the reactants are held together in a defined geometry and at an internuclear separation which corresponds to a minimum value in the interaction potential. In the present work the reaction is initiated by absorption of a photon and subsequently takes place with a restricted range of impact parameters governed by a thermal distribution of molecular displacements about the equilibrium geometry. In addition, the reactive encounters are abbreviated owing to the fact that reaction begins at a configuration which lies well into the entrance channel of the potential-energy surface (PES), i.e. in the transition state (TS) region.

In the photoinduced reactions of molecules ordered at surfaces, photolysis of an adsorbed molecule generates a recoiling fragment which can then react with an adjacent molecule co-adsorbed on the surface. This is referred to as Surface Aligned Photochemistry[1]; examples studied in this laboratory include the reactions $\vec{\text{H}} + \text{H}_2\text{S}(\text{ad}) \rightarrow \text{H}_2(\text{g})$ [3], $\vec{\text{S}} + \text{SCO}(\text{ad}) \rightarrow \text{S}_2(\text{g})$ [4], and $\vec{\text{H}} + \text{HBr}(\text{ad}) \rightarrow \text{H}_2(\text{g})$ [5] in which the reactant was adsorbed on a LiF(001) surface. The dynamics can be inferred from the motion of products leaving the surface, measured using such techniques as time-of-flight mass spectrometry with angular resolution.

The use of photoinduced reaction in van der Waals complexes to obtain information on reaction dynamics was pioneered by Wittig et al.[2] and by Soep and co-workers[6-11]. A brief summary of the features of their systems is given in the following paragraphs.

In Wittig's group, photodissociation of one of the constituents in a binary-molecular complex liberated a reactant which recoiled toward the other constituent. For instance, photolysis of the hydride, HX, within the complex, $\text{CO}_2 \cdots \text{HX}$, liberated H-atoms which proceeded to react with the CO_2 molecule at a preferred angle defined by the geometry of the precursor complex. The translational energy of the incident H-atom was related to the photon energy minus the dissociation energy for the hydride molecule.

The probability of reaction was found to be dependent on the structure of the complex. For example, the structure of $\text{CO}_2 \cdots \text{HX}$ is linear for $\text{X}=\text{Cl}$, while the geometry of the constituents for $\text{X}=\text{Br}$ is bent with the Br near the C atom. The yield of OH produced from $\text{CO}_2 \cdots \text{HBr}$ photolysis was found to be $\sim 40\times$ more than from $\text{CO}_2 \cdots \text{HCl}$ photolysis at similar values for the initial H-atom translational energy. This indicated that, in some cases, the geometry of the constituents aligned in a complex could be favorable for reaction to occur and in other cases less so.

In addition, when more than one product channel existed, different branching ratios were sometimes observed compared to those for the corresponding gas-phase reaction. The photoinitiated reaction in $\text{N}_2\text{O} \cdots \text{HX}$ ($\text{X}=\text{Cl}, \text{Br}$) has two product channels, $\text{OH}(\text{X}_2\text{P}) + \text{N}_2(\text{X}^1\text{S})$ and

$\text{NH}(\text{X}^3\text{S}) + \text{NO}(\text{X}^2\text{P})$. For photolysis of $\text{N}_2\text{O}\cdots\text{HI}$, the $[\text{NH}]/[\text{OH}]$ ratio was reduced by an order of magnitude relative to its value from photolysis of (uncomplexed) gaseous $\text{N}_2\text{O}/\text{HI}$ mixtures. This observation is related to the minimum-energy structure for this complex, in which the H atom in the HI molecule is located adjacent to the O atom in the N_2O molecule, leading to a preference for the $\text{OH}+\text{N}_2$ product channel.

In Soep's group, complexes of metal atoms with inorganic molecules, in particular $\text{Hg}\cdots\text{Cl}_2$ [6], $\text{Hg}\cdots\text{H}_2$ [7] and $\text{Ca}\cdots\text{XH}$ (where $\text{X}=\text{F}$ [11], Cl [8-11], Br [9-11] and I [11]) were studied. Photoexcitation of the metal-atom chromophore within the complex accessed electronically-excited states of the complex from which reaction could then occur. The electronic states prepared by photoexcitation of the complexes correlated with the $\text{Hg}(^3\text{P}_1)$ and $\text{Ca}(^1\text{P}_1)$ states in the asymptotic limit. In the environment of the complex the atomic states are split into components which correspond to Σ and Π configurations of the p-orbital relative to the closed-shell molecule.

State-selective photoexcitation of the metal atom in the complex allowed analysis of the effect of the orientation of the p-orbital relative to the internuclear axis on the reactivity and the product-state distribution. This was achieved by varying the wavelength used to excite the ground-state complex and either monitoring emission from electronically-excited states of the product, or probing the electronic-ground state by laser-induced fluorescence (LIF). The appropriate signal-intensity measured as a function of the excitation wavelength was referred to as an 'action spectrum' for the system.

The excited-state reaction of $\text{Hg}\cdot\cdot\text{H}_2$ was observed to depend on the entrance-channel orbital orientation since the Hg atom retains an electron in a p orbital during the course of the reaction to give $\text{HgH}(\text{X } ^2\Sigma) + \text{H}$. The action spectrum identified two channels for reaction; the first occurring rapidly through the Π -configuration of the metal p-orbital and the second occurring more slowly through the Σ -configuration. Product analysis indicated a greater product-rotational excitation for reaction through the Π -configuration.

In contrast, selectivity for the product channel was not observed with $\text{Hg}\cdot\cdot\text{Cl}_2$ and $\text{Ca}\cdot\cdot\text{ClH}$ complexes, even though state-selective preparation of the excited complexes was achieved. For example, in the action-spectrum for $\text{Ca}\cdot\cdot\text{ClH}$ the intermolecular bending mode was observed as a vibrational progression. In this case, the excited complex evolved into a configuration in which the covalent and ionic, (i.e. $\text{Hg}^+\cdot\cdot\text{Cl}_2^-$ or $\text{Ca}^+\cdot\cdot\text{ClH}^-$) states were degenerate. The result was charge-transfer in the excited complex in which the metal p-orbital electron was transferred to the halide. Thus, the excited-state reactions of $\text{Hg}\cdot\cdot\text{Cl}_2$ and $\text{Ca}\cdot\cdot\text{ClH}$ were not observed to have specificity with respect to orbital orientation, since the Hg and Ca atoms lose the p-orbital electron early in the course of the reaction.

The experiments performed by Wittig et al. and Soep and co-workers have been described as Transition State Spectroscopy[12]. For instance, photodissociation of the hydride, HX , within the complex $\text{CO}_2\cdot\cdot\text{HX}$, forms an excited species, $[\text{O}-\text{CO}\cdot\cdot\text{H}]^\ddagger$, at a configuration intermediate between the reactants, CO_2+H (at infinite O-H separation) and products, $\text{CO}+\text{OH}$ (at infinite C-O separation), for the reaction

$\text{CO}_2 + \text{H} \rightarrow \text{CO} + \text{OH}$. This excited species, indicated by the double-dagger symbol, is the Transition State (TS) for the above reaction. Similarly, photoexcitation of the complex, $\text{Ca} \cdots \text{XH}$, forms a TS, $[\text{Ca}^* \cdots \text{XH}]^\ddagger$ for the excited-state reaction $\text{Ca}^* + \text{XH} \rightarrow \text{CaX} + \text{H}$. The initial configuration and internal energy of the TS in both these examples will depend on the excitation wavelength. By varying the wavelength used to form the excited state, the dynamics can be probed as a function of different starting configurations in the TS region. The possibility of achieving a spectroscopy of the Transition State was first realized in this laboratory[12]. Relevant advances in this field will be described in the next section. It will be evident from the foregoing that the TS is defined as any configuration intermediate between reagents and products (rather than a limited set of configurations critical to the reaction rate, as is sometimes done).

The object of this thesis was to develop a Transition State Spectroscopy for the classic charge-transfer reactions of the type $\text{Na}^* + \text{XR} \rightarrow \text{NaX} + \text{R}$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$, $\text{R}=\text{CH}_3, \text{Ph}, \text{H}$), using the technique of photoinitiated reaction in van der Waals complexes. Tunable monochromatic excitation of the stable ground-state complex, $\text{Na} \cdots \text{XR}$, accessed the PES for the excited-state reaction at a selected configuration intermediate between reagents ($\text{Na}^* + \text{XR}$) and products ($\text{NaX} + \text{R}$). Charge-transfer then occurred in a narrow region of configurations of the evolving TS, followed by dissociation of XR^- to give the product NaX molecule. This process is symbolized in eq. 1.1.

The charge-transfer process is often referred to as 'harpooning'[13-15] and occurs when the ionic and covalent configurations of the excited complex are degenerate. In this work, the excitation wavelength λ_1 (corresponding to ν_1) was varied to access the TS in a series of different starting configurations.

Previous work performed in this group focused on the photoinduced charge-transfer dissociation reaction in $\text{Na}_2\cdots\text{ClCH}_3$ [16,17], $\text{Na}\cdots\text{FPh}$ [17] and $\text{Na}\cdots(\text{FCH}_3)_n$ ($n=1-4$)[18,19] complexes. The present work presents a comparative study of the photoinitiated reaction in $\text{Na}\cdots\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) complexes as well as a detailed study of the dynamics of the reaction $\text{Na}^* + \text{HF} \rightarrow \text{NaF} + \text{H}$, in which the experimental photodepletion spectrum for the complex $\text{Na}\cdots\text{FH}$ is compared with an *ab initio* computation (performed by others in this laboratory) of the PESs for the $\text{Na}\cdots\text{FH}$ and $\text{Na}^*\cdots\text{FH}$ systems.

The study of van der Waals clusters could at a future date serve to bridge the gap between the gas-phase and surface photoreactions. This would be achieved by systematically increasing the size of the metal component in the cluster. Photoinduced dissociation of small molecules on surfaces (including surfaces containing alkali metals) has also been studied in this laboratory[20-22]. Electronic excitation of a substrate surface can lead to either electronic-energy transfer to the adsorbate (e.g. from $\text{LiF}(001)$ to OCS [20]) or charge-transfer to the adsorbate (e.g. from $\text{Ag}(111)$ to CCl_4 [21]) resulting in either case in bond cleavage. The latter process has also been observed to trigger a reaction between the adsorbate and the surface in the series of chloromethanes, CCl_4 , CHCl_3 , CH_2Cl_2 , and CH_3Cl [22],

The negative ion, RCl^- , formed by way of charge-transfer from the substrate was shown to react with the surface prior to R-Cl bond-cleavage. In addition, a pathway (referred to as autoionization of RCl^-) which would correspond to a failed reaction, in which charge-transfer led to desorption of RCl with the electron being transferred back to the surface, has also been identified[22].

1.2 Transition State Spectroscopy

In this work, as noted above, we define the Transition State (TS) to include all the configurations of the system intermediate between reactants and products[12]. The field of Transition State Spectroscopy (for a recent review see ref. 23) began with experimental observation of the TS in the emission and absorption spectra of reacting systems (for a more detailed review of these experiments see ref. 24). Recent developments have seen the evolution of the TS probed in real-time using ultrafast-laser pulses[25].

In the previous section, formation of a TS in a well defined configuration by monochromatic excitation of a stable van der Waals complex was described. Imre, Kinsey and co-workers have achieved a similar result for photodissociation by laser excitation of a bound molecule. The excitation accessed an electronically-excited state which was repulsive along one coordinate. Emission to quantum states of the electronic-ground state occurred as the TS dissociated into products. A study of the photodissociation 'half reactions' $[\text{CH}_3\text{I}]^\ddagger \rightarrow \text{CH}_3 + \text{I}$ [26] and $[\text{O}_3]^\ddagger \rightarrow \text{O}_2 + \text{O}$ [27] was made using this technique. The emission spectrum

for these systems contained information regarding the ground and excited-state PESs and the dissociation dynamics on the excited PES.

Negative-ion photodetachment has provided an additional technique for the preparation of the TS in a defined configuration from a bound precursor. The dynamics of reactions $A + BC \rightarrow AB + C$ was probed by photoelectron spectroscopy of the stable negative-ion ABC^- . This approach, due to Neumark and co-workers, has provided detail on the TS for such important reactions as $F + H_2 \rightarrow HF + H$ [28,29]. The translational energy distribution of the ejected electrons from $HHF^- \rightarrow [HHF]^\ddagger + e^-$ contained information on vibrational modes approximately orthogonal to the reaction coordinate, and also quantum "bottlenecks" along the reaction pathway.

The study of the evolution of the TS in the time-domain was pioneered by Zewail and co-workers[25]. In these experiments the TS was prepared in a defined geometry and the motion of the wavepacket was probed using ultrafast lasers. One system studied by this group has relevance to the work in this thesis. The dynamical motion of a wavepacket along the crossing region of covalent and ionic potential-energy curves was examined for the reaction $NaI^* \rightarrow [Na \cdots I]^\ddagger \rightarrow Na + I$. The time-dependent formation of product Na atoms from the excited-state dissociation reaction was probed following excitation of the ionic-ground state of NaI to the covalent-excited state, NaI^* . The observation of an oscillation of the Na signal on the femtosecond time-scale indicated that some NaI^* molecules were being trapped in a well formed by the avoided crossing of the covalent and ionic PES. It is in this region of configurations in which charge-transfer harpooning occurs in the

reaction $\text{Na} + \text{I} \rightarrow \text{NaI}$. This femtosecond Transition State Spectroscopy gave information on the traversal time of the wavepacket across the well and the probability of escape into the channel leading to the products, $\text{Na} + \text{I}$ [30].

1.3 Harpooning reactions

The harpooning mechanism was proposed over 60 years ago by M. Polanyi to account for the cross-sections observed in the reaction of alkali metals with halogen molecules[13]. The experimentally measured cross-section for these reactions were much larger than predicted by simple collision theory, even if reaction was assumed to occur at every atom+molecule collision. Charge-transfer between the alkali metal, M, and the halogen, X_2 , was proposed to occur at a distance much greater than the normal collision diameter, where the ionic and covalent states of $\text{M} \cdots \text{X}_2$ are degenerate. The charge transfer yields X_2^- which dissociates as $\text{M}^+ \text{X}^-$ is formed. The basic features of this mechanism were developed in a theoretical study by Magee[14] and are illustrated in fig. 1.1. This model provided some good qualitative results. The cross-section for the harpooning reactions were observed to increase with decreasing ionization energy of the metal, $I^\circ(\text{M})$; in the order $\text{M} = \text{Li} < \text{Na} < \text{K} < \text{Rb} < \text{Cs}$ and $\text{Na}(3s^2S) < \text{Na}(3p^2P) < \text{Na}(4s^2S) < \text{Na}(3d^2D)$, and with increasing electron affinity of the halogen, $E^\circ(\text{X}_2)$; in the order $\text{X} = \text{I} < \text{Br} < \text{Cl} < \text{F}$ [15].

Harpooning reactions have since been studied in detail in molecular-beam experiments, particular by Herschbach and co-workers[15,31-36]. A stripping mechanism was found for reactions which have a large cross-section, such as those between alkali metals and

$$-e^2/R_c + \Delta E_0^0 = -C/R_c^6 \quad (\approx 0)$$

$$R_c \approx e^2/\Delta E_0^0$$

$$\sigma \approx \pi R_c^2$$

Figure 1.1: The covalent, $M+X_2$, and ionic, $M^++X_2^-$, interaction potentials for the alkali metal and halogen molecule. The curves intersect at an internuclear distance, R_c , at which the covalent potential is described (approximately) by $-C/R^6$ and the ionic potential is described by $-e^2/R+\Delta E_0$, relative to the $M+X_2$ asymptotic limit.

halogen molecules. Angular distributions indicated that the products recoiled largely into the forward hemisphere. This anisotropy suggested a direct or impulsive mechanism in which the duration of the reactive collision was well below the average rotational period for a collision complex. Kinematic analysis showed that the final relative translational energy of the products was comparable to the initial kinetic energy of the reactants, so that most of the substantial energy released must appear as internal excitation of the products. Electric-deflection experiments demonstrated that only a small part of this internal energy appeared in rotation, indicating that most of the reaction exoergicity was disposed in vibrational modes. This is consistent with an early barrier on the potential-energy surface (PES)[37], expected for highly exoergic reactions.

In contrast, the reaction of alkali-metal atoms with methyl-halide and hydrogen-halide molecules are characterized by a much lower cross-section. A larger flux for the reaction products are observed in the backward hemisphere of the angular distribution. Many of these reactions are described by a late barrier on the PES. The characteristics of these reactions will be described in more detail.

Reaction of Na atoms with methyl halide molecules

The charge-transfer reactions $\text{Na} + \text{CH}_3\text{X} \rightarrow \text{NaX} + \text{CH}_3$ ($\text{X} = \text{F}, \text{Cl}, \text{Br}$) are exoergic but possess a significant energy barrier. The origin of the energy barrier for the ground-state reaction is the low electron affinity of the halide molecule and the corresponding shorter internuclear distance at which electron transfer can then occur. For $\text{Na} + \text{CH}_3\text{X}$, the electron transfer is thought to take place at a distance close to the equilibrium

bond length in Na-X[38] and it is postulated that CH₃-X must be stretched prior to charge-transfer[17,19]. This is in accord with experimental observation in molecular beams of a very anisotropic angular distribution for the reaction products, in which scattering occurs predominantly in the backward hemisphere. For example, the system K+ICH₃ was shown by Bernstein and co-workers to proceed by a rebound mechanism[39]. The total integrated cross-section for this reaction was lower than for the corresponding reaction with the halogen molecule, I₂.

The orientation of methyl-halide molecules in molecular beams using electric fields have indicated a steric effect in which the reaction cross section is much lower if the metal atom approaches from the methyl end of the halide molecule. This was shown by Brooks and co-worker for K+ICH₃[40] and by Bernstein et al. for Rb+ICH₃[41]. This is also consistent with the requirement for short Na-X distances at which charge-transfer can occur.

The presence of a shallow potential-energy well ahead of the large energy barrier in the ground state enables the formation of complexes between Na and XCH₃ under conditions of low collision energy and in the presence of a third body to remove the excess internal energy. The geometry of the van der Waals complexes will have a large effect on the outcome following photoexcitation (see experiments by Wittig[2]). Trenary et al.[42] have conducted *ab initio* calculations on a family of molecular complexes involving Li and Na atoms with dipolar molecules such as NH₃, H₂O, HF, PH₃, H₂S and HCl. Their calculations indicate that the minimum-energy structures had the negatively-charged end of the molecular component, such as the halogen X when present, adjacent

to the alkali atom. Thus, the van der Waals complexes, $\text{Na}\cdots\text{XCH}_3$, are favorably oriented for charge-transfer dissociation to occur.

Translational energy thresholds for the reactions of alkali metals with methyl halides were determined by Bernstein and co-workers for the reactions of K and Rb with CH_3Br and CH_3I [43-45]. The reaction cross section was observed to increase with the translation energy, E_{tr} , just above threshold but reaches a peak which is followed by a decline to a minimum with increasing E_{tr} . The maximum in the cross-section, at an intermediate collision energy, was also noted in the reaction of Na with CH_3I by Ureña et al.[46]. The initial increase in the reaction cross section with collision energy was explained by Herschbach and co-workers[47] as arising from the existence of an energy barrier in the entrance channel of the PES due to the avoided crossing between the covalent (MXCH_3) and ionic (M^+XCH_3^-) states. The subsequent decrease arises from molecular trajectories which cross back from the ionic to the covalent state without becoming trapped.

The energy barriers have been estimated for the reactions of alkali atoms with various methyl-halide molecules by considering a simple electron-jump model using semi-empirical potential-energy curves for the CH_3X molecule and CH_3X^- ion[48]. The barrier for the reaction $\text{M}+\text{CH}_3\text{X}$ was described as the energy required to stretch the C-X bond in order to obtain a positive electron affinity for the halide molecule. The values obtained were 0.026eV for CH_3I , 0.25eV for CH_3Br , 0.54eV for CH_3Cl , and 1.90eV for CH_3F . The former two results were noted by Wu in ref. 48 to be in very good agreement with experimental measurements by Bernstein and co-workers[43-45] of the translational energy thresholds

for the reactions of K/Rb with CH_3I and CH_3Br . However, a similar comparison was not made in ref. 48 with experimental values for ClCH_3 and FCH_3 because the collision energies obtainable in Bernstein's experiments were limited to values below 0.5eV and products from the reaction of alkali metals with FCH_3 or ClCH_3 were not detected.

A cross-section of $6\text{\AA}$ for the reaction of $\text{Na}(3s\ ^2\text{S})$ with BrCH_3 was measured by Y.T. Lee and co-workers using a collision energy of 0.91eV[49]. The reaction of ground and excited-state Na with ClCH_3 was found to be too small in ref. 49 for accurate measurements of the cross-section and angular distribution of the reaction products. In addition, measurements of the energy threshold for reaction in molecular-beam experiments have not been made. However, sodium-flame experiments for $\text{K} + \text{ClCH}_3$ systems enabled an estimate of the activation energy for $\text{Na} + \text{ClCH}_3$ of 0.36 eV[50]. This should be compared with the calculation using the electron-jump model described above which estimated an energy barrier of 0.54 eV. A considerably more accurate *ab initio* calculation using single-reference coupled-cluster theory with single, double and triple-excited components gave a value of 1.56eV[51] for the energy required to stretch the F- CH_3 bond (in this case by $\sim 1.4 \times r_0(\text{F}-\text{CH}_3)$) in order to obtain a positive electron affinity. This is lower than the energy barrier determined by Wu in ref. 48.

The reaction of excited-state Na^* with methyl bromide has been studied by Lee and co-workers[49]. Although the reaction still proceeds by the rebound mechanism, with increasing electronic excitation the reactive cross section was observed to increase and the width of the center-of-mass angular distribution increased. The steric angle appears

to be widened with increasing electronic energy and, thus, the metal-atom valence-orbital size[49].

Reaction of Na atoms with HF molecules

The reaction of alkali-metal atoms with hydrogen-halide molecules were not studied in as much detail in molecular beams until more recently due to unfavourable kinematics in the center-of-mass distribution for these systems. However, experimental studies have now been performed by, among others, Brooks[52,53], Loesch[54-60], Lee[61-64], and Düren[65-68] and their co-workers. The reaction $\text{Na} + \text{HF} \rightarrow \text{NaF} + \text{H}$ is endoergic by 30kcal/mol. The dependence of the cross-section for the ground-state reaction on vibrational and rotational excitation of the HF molecule was examined earlier in this laboratory by infrared-chemiluminescence depletion[70,71]. Reaction was not observed with HF molecules in the ground state, however, vibrational excitation gave a large enhancement of cross-section[71] consistent with a 'late barrier'[37].

In contrast, in work also performed in this laboratory, the reaction cross-section was observed to initially decrease as the reagent rotational energy was increased[70]. This was ascribed to rotation of the HF molecule out of the favored orientation for reaction to occur[60,70] indicating the existence of a steric effect similar to that observed in the reaction of alkali metals with methyl-halide molecules. However, at large values of the rotational-quantum number, the reaction cross section did increase due to vibrational stretching modes which were induced by high

energy rotational motion, resulting in weakening of the H-F bond under attack.

The reaction of excited-state Na^* with HCl has been extensively studied by Lee and co-workers using crossed-molecular beams[62-64]. They observed an increasing reactive cross-section and decreasing product recoil energy with increasing electronic energy of Na^* . This observation is consistent with a harpooning model for the reaction. The reaction dynamics were thought to change with electronic excitation. Excited Na^* has a lower ionization potential and hence gives rise to an earlier electron transfer. However, for the case $\text{Na} + \text{HF}$, they observed no reaction of Na^* even at large electronic excitations and high collision energies.

The model 'Direct Interaction with Product Repulsion' (DIPR)[72,73] has been used successfully to simulate the dynamics in harpooning reactions which are characterized by high cross-sections. The basis of this model is the assumption of sequential release of repulsive and attractive energy in which the interaction force decreases monotonically with time. A discrepancy between the experimental observations and DIPR simulations for the systems $\text{Na}^* + \text{HCl}$, in which Na is in both ground and excited states, was noted by Lee's group[62]. The main assumption in this case breaks down and the reaction possesses a greater degree of concertedness.

Reactive scattering of $\text{Na}^*(3p\ ^2P)$ from HF was observed in crossed-molecular beams by Düren and co-workers[66-69], with a significant reaction cross section of $1\text{-}2\text{\AA}^2$. The absence of reactive signal in the

experiments from Lee's group was due to the lower sensitivity of their detection method[65]. From the rainbow scattering in non-reactive collisions, estimates were obtained for the well-depths in the entrance valley of the PESs for the ground and excited state. For the electronic-ground state they obtained a well depth of 65meV for approach of the Na atom to the F-end of the halide molecule. However, this value will represent an average across all Na+HF collision geometries and, therefore, a lower limit of the actual well depth for the equilibrium geometry of the complex.

In the present study of TSS the experimental findings have been supplemented by *ab initio* calculation of the ground and excited-state PES. The ground-state PES of the Na $\cdot$ FH system has been analyzed before using semi-empirical[74] and *ab initio*[75-78] approaches.

The semi-empirical valence-bond calculation due to Shapiro and Zeiri[74] indicated that the saddle point on the ground-state PES, which corresponds to the minimal-energy barrier, occurred in a non-linear geometry at an angle of $\sim 60^\circ$ and a height of 29.2kcal/mol. It was in this region where charge-transfer between covalent, NaFH, and ionic, Na $^+$ FH $^-$, states was believed to occur. The explanation given by Shapiro and Zeiri[74] for the approach of the Na atom in the non-linear geometry was due to stabilizing the second ionic structure Na $^+$ +H $^-$ F which would lower the (HF) $^-$ repulsion due to the possibility of H-F $\leftrightarrow$ HF $^-$ resonance. The saddle point was located in the exit channel in which the HF molecule was separated twice as much as NaF from its equilibrium position.

The origin of the barrier in the exit channel has already been described as due to stabilization of the ionic configuration by the close proximity of Na^+ and F^- , with the H atom separated significantly from F^- [38]. A considerably more accurate *ab initio* calculation using single-reference coupled-cluster theory with single, double and triple-excited components gave a value of 21.3 kcal/mol for the energy required to stretch the F-H bond (in this case by $1.28 \times r_0(\text{H-F})$) in order to obtain a positive electron affinity such that charge-transfer from the alkali metal to HF can occur [51].

Experimental evidence that a non-linear geometry for the transition state would be favored in the ground state reactions was provided by Herschbach and co-workers in their crossed-molecular beam studies of the reverse process, $\text{H} + \text{NaF} \rightarrow \text{Na} + \text{HF}$ [79]. The differential reaction cross-section was sideways peaked in the center-of-mass angular distribution with respect to the incident H atoms, indicating a likely preference for non-linear reaction geometries.

The restricted Hartree-Fock *ab initio* calculation by Paniagua and co-workers only considered collinear-reaction geometries for $\text{Na} + \text{HF}$ [75]. Consequently, a much larger value of 40.9 kcal/mol was reported for the barrier height in the ground-state reaction. The barrier was again found to be located in the exit channel at which the Na-F bond length was nearly at its equilibrium value and the H-F bond was considerably stretched ($1.54 \times r_0(\text{H-F})$). This configuration lies much deeper into the exit channel than calculated by Shapiro and Zeiri for the bent geometry of NaFH. However, this calculation also identified a bound state of NaFH with a potential-energy well of 2.5 kcal/mol which could stabilize an

intermediate complex in the entrance channel of the ground-state PES. The potential-energy minimum occurred at a stretched Na-F distance relative to the equilibrium bond length in the NaF molecule ($1.33 \times r_0(\text{Na-F})$) and at a slightly extended H-F distance ($1.06 \times r_0(\text{H-F})$). Earlier Trenary et al. had calculated at the SCF level an interaction energy of 1.2 kcal/mol between Na and HF[42]. The atoms were fixed in a linear geometry and only the Na-F distance was optimized (in this calculation at $1.27 \times r_0(\text{Na-F})$). However, this energy was noted to represent a lower limit due to the neglect of configuration interaction in their calculation.

Laganá et al. performed an extended *ab initio* calculation at a series of geometries for the system; 180° , 120° , 90° , 75° , 60° and 45° [77]. They identified a barrier for the ground-state reaction in the exit channel of the PES closest to the 75° geometry and at an energy of 20.5 kcal/mol above the reactant asymptote. The Na-F distance was almost at the equilibrium bond length in the product molecule and the H-F distance was greatly extended ($1.78 \times r_0(\text{H-F})$). Compared with Shapiro and Zeiri's semi-empirical calculation, this H-F distance was slightly longer and the minimum-energy geometry was at a slightly wider angle. In addition, this calculation also identified a potential-energy well in the entrance channel of the PES. The geometry of the ground-state complex was determined to be non-linear with a bond angle between 75° and 90° , in which the Na-F bond was slightly extended ($1.16 \times r_0(\text{Na-F})$). The depth of the well was reported to be 11.4 kcal/mol; however, this value appears to have been overestimated in their calculation.

Quantum-classical trajectory calculations on the PESs determined by Laganá and co-workers[78] enabled a fairly accurate simulation of the

experimental results obtained for the vibrational and rotational dependence of the reaction $\text{Na} + \text{HF}$ in refs. 70 and 71. However, the authors of this theoretical work did suggest that the depletion was largely associated with rotationally inelastic non-reactive collisions. Their calculations indicated that the cross-section would increase with vibrational excitation and decrease sharply with increasing collision energy. This was explained due to an attractive interaction at long range in which the early location of the well stabilizes an intermediate complex. The potential-energy well is less capable of capturing large impact-parameter trajectories when the collision energy increases. However, an increase in impact parameter does occur as the vibrational energy increases. The fraction of rotational energy in the product was found to be fairly independent of initial vibrational and collision energy. The initial vibrational energy in the reactant was found to be transferred to the product in the course of the reactive encounter.

The theoretical information about excited states has hitherto been limited to a small-scale calculation of the valence bond and configuration-interaction type[76]. On the basis of the shapes of the adiabatic PESs, the authors of this theoretical study suggested that the reaction of Na^* with HF was associated with surface hopping from the first excited state to the ground state PES into the product channel $\text{NaF} + \text{H}$. Surface hopping mechanisms have been proposed for other systems of electronically-excited alkali metal atoms reacting with hydrogen halide molecules; i.e. $\text{Li}^* + \text{HF}$ [80], $\text{Na}^* + \text{HCl}$ [81,82], $\text{Li}^* + \text{HCl}$ [83].

The surface hopping mechanism will be discussed in this thesis as a likely description of the dynamics in the photoinitiated reaction of

Na $\cdots$ FH complexes. Excitation of the stable complex, Na $\cdots$ FH, on the ground-state PES would access the transition state (TS) for the excited-state reaction in a configuration in which the Na-F bond length is extended relative to its equilibrium value and the H-F bond length is nearly at equilibrium. The transition state would then evolve into a configuration in which the ground and excited state PES approach each other closely. This would be located above the saddle point of the ground-state PES in which the Na-F bond length is compressed and the H-F bond length significantly extended. The product channel, NaF+H, would then be accessed by surface hopping from the excited state.

Recently, on the basis of *ab initio* studies of the systems Li + HF[84], Na + HCl[82] and a semi-empirical study of the system Li + HCl[85], laser catalyzed reaction schemes were proposed. This mechanism described excitation of a collision complex, M $\cdots$ XH, followed by stimulated emission from the first excited state into the product channel on the ground state.

1.4 Summary and thesis preview

The reactions of alkali metals in the ground and electronically-excited states with halogen and halide molecules have received considerable attention. Charge-transfer from the alkali-metal atom to the halide is thought to occur in the Transition State (TS) across a narrow range of configurations. This is the 'harpooning' event, first proposed over half a century ago to be the crucial step in these reactions. The negatively-charged halide molecule, XR $^-$, then dissociates to yield the product, NaX.

Photoinitiated reaction in the van der Waals complexes, $\text{Na}\cdots\text{XR}$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$, $\text{R}=\text{CH}_3$, and $\text{X}=\text{F}, \text{R}=\text{H}$), provides a novel approach to the study of the dynamics of these harpooning reactions by way of Transition State Spectroscopy. Photoexcitation of $\text{Na}\cdots\text{XR}$ accesses a potential-energy surface (PES), which would be traversed by the excited-state reaction, $\text{Na}^*+\text{XR}\rightarrow\text{NaX}+\text{R}$, at a selected configuration intermediate between the reactants, Na^*+XR , and the products, $\text{NaX}+\text{R}$, i.e. a selected Transition State, TS; $[\text{Na}^*\cdots\text{XR}]^\ddagger$. The starting configuration for the TS is determined by the geometry of $\text{Na}\cdots\text{XR}$ and the excitation energy, $h\nu_1$. Therefore, by varying the excitation wavelength, λ_1 , it should be possible to probe the excited-state dynamics as a function of selected TS configurations.

The large energy barrier for the ground-state reaction of Na with HF and XCH_3 ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) enables the formation of complexes under conditions of low collision energies and in the presence of a third body to remove the excess internal energy. The presence of a potential-energy well on the ground-state surface has been predicted by theoretical calculations on the system NaFH [42,72,75].

A comparative study of the excited-state-harpooning reactions between Na and XCH_3 ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) has been made in this work. The ground-state complexes were excited to PESs traversed by the reaction of excited-state Na^* with XCH_3 . By increasing the excitation energy we were able to access the TS for the reaction with successively higher electronically-excited states of Na^* ; including $3p\ ^2P$, $4s\ ^2S$ and $3d\ ^2D$. The technique used to probe the dynamics of the excited states was to

monitor depletion of the ground-state complex as a function of the excitation wavelength.

The dynamics of the excited-state reaction of $\text{Na}^*(3p\ ^2P)$ with HF will be discussed in greater detail. The increased simplicity obtained by substituting a H atom for a methyl group has enabled extensive *ab initio* calculations to be performed on this system (by others in this laboratory) which make a more detailed description of the reaction dynamics possible. Dissociation of the TS has been suggested to occur via surface hopping from the electronically-excited state into the product channel on the ground-state PES[76]. The region of configurations in which the character of the TS abruptly changes from largely covalent to largely ionic, $[\text{Na}^+\cdots\text{FH}]^\ddagger \rightarrow [\text{Na}^+\cdots\text{FH}^-]^\ddagger$ will be explored.

Another approach to Transition State spectroscopy will be described for the reaction $\text{Na}^* + \text{FCH}_3 \rightarrow \text{NaF} + \text{CH}_3$. In this case, the stable ground-state complex, $\text{Na}\cdots\text{FCH}_3$, was excited to the PES for the excited-state reaction and then the decay of the TS, $[\text{Na}^+\cdots\text{FCH}_3]^\ddagger$, was probed directly by photoionization from the TS. This experiment, performed in the frequency-domain, was able to provide an order-of-magnitude estimate of the lifetime for the TS in the harpooning reaction.

Chapter 2.**Experiment**

The van der Waals complexes, $\text{Na}\cdots\text{XR}$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}, \text{R}=\text{CH}_3$; $\text{X}=\text{F}, \text{R}=\text{H}$) were formed using the crossed-molecular beam technique pioneered by Hertel and co-workers[86]. A Na beam, generated from an oven source, was directed into the expansion region of a pulsed-supersonic jet of the appropriate halide in an inert-carrier gas. The concentration of complexes was measured a short distance away from the crossing region of the molecular beams by photoionization time-of-flight (TOF) mass spectrometry. Reaction was found to be initiated by photoexcitation of the complexes across a broad range of wavelengths. The complex-signal intensity was measured in the presence and absence of a single excitation wavelength in order to estimate the fraction of complexes depleted as a function of the wavelength. A schematic diagram of the experimental apparatus is given in fig. 2.1. The apparatus has been modified from that used in earlier experiments from this laboratory (see refs. 16 to 19). The distance of the crossing point for the two molecular beams relative to each source has been significantly reduced, along with the distance from the crossing point to the extraction region of the time-of-flight mass spectrometer (TOFMS). This was found to give a greater yield of the weakly-bound complexes for accurate depletion measurements. Additional ion optics were also installed at the early stages of the flight path in order to increase the detection efficiency for the complex ions. These modification will be described in detail below.

Figure 2.1: Side view of the vacuum system

2.1 Experimental apparatus

Vacuum system

The vacuum system comprised three differentially-pumped chambers, A, B and C (see fig. 2.1). Chamber A contained the molecular-beam sources (see below) and the extraction region for the TOFMS. The excitation laser, L_1 , and the ionization laser, L_2 , were counterpropagated through chamber A, perpendicular to the axis of the TOFMS. This chamber was pumped by a 10" oil-diffusion pump (Varian VHS 10). The diffusion pump was connected to a two-stage rotary backing pump (Edwards E2M40). The second acceleration stage of the TOFMS was located in chamber B which was pumped by a 4" oil-diffusion pump (NRC VHS 4). The 10" and 4" diffusion pumps were baffled by liquid-nitrogen traps (Varian 0362 Cryo baffles with NRC882 LN₂ controller). The ion signals were measured in chamber C which was pumped by a 2" oil-diffusion pump (Edwards diffstak series 63). The 4" and 2" diffusion pumps were both backed by single-stage rotary backing pumps (Edwards E1M5). The average background pressures under typical operating conditions were 2×10^{-5} torr, $< 10^{-6}$ torr and $< 10^{-7}$ torr respectively, for chambers A, B, and C. The pressure in each chamber was measured by a Bayard-Alpert ionization gauge (Kurt J. Lesker model 4336, with Granville-Phillips controller, series 260).

Crossed-molecular beam assembly

The van der Waals complexes were formed by crossing a Na beam generated by a stainless-steel oven with the expansion region of a pulsed-supersonic jet of the appropriate halide (XCH_3 , $X=F$, Cl, Br, or

FH) in an inert-carrier gas (He or Ar). This arrangement is illustrated in fig. 2.2. The oven consisted of a cylindrical reservoir connected to an elbow-shaped nozzle extension. The construction of the oven has been described in detail earlier[87]. The atomic beam emerged from a 0.8mm-diameter orifice along a 10mm-length channel in the nozzle section. The reservoir and nozzle were independently heated by means of custom designed, porous, ceramic-heating elements, embedded with Fe/Cr/Ni alloy wires (Thermcraft). The temperature of each section was measured with K-type thermocouples (Omega Eng., Inc.) and monitored by means of two separate temperature controllers (Omega Eng., Inc., CN2002K-A-PID2). The reservoir was heated to a temperature of 490°C, while the nozzle section was heated slightly higher, to a temperature, T , of 520°C in order to avoid condensation of Na in the exit channel. These temperatures correspond to a sodium pressure, p , of ~ 8 torr ($\sim 10^{17}$ particles/cm³) ahead of the nozzle, with an equilibrium composition of approx. 94% Na, 6% Na₂[88]. However, following expansion from the orifice, the atomic beam will likely contain an increased proportion of Na₂ molecules[89,90]. The mean-free path of a gas at low pressure is given by $\lambda = kT / (2^{1/2} \pi r^2 p)$ where r is the radius of the metal atom. This gives a value of $\lambda \approx 4 \times 10^{-3}$ mm for Na atoms at the nozzle exit (using a metal atom radius of $r = 2.27 \text{ \AA}$ [91]). The atomic beam was characterized by a Knudsen number (nozzle diameter/ mean free path) of ~ 200 , which corresponds to a well-developed supersonic expansion[92]. The final kinetic energy of an expanded-supersonic beam is $(5/2)kT$ which corresponds to a terminal velocity of ~ 1200 m/s for the Na atoms in the atomic beam.

Figure 2.2: Crossed-molecular beam assembly- 1, stainless-steel oven in contact with ceramic heating elements; 2, water-cooled jacket; 3, Na-beam collimator; 4, pulsed valve.

The oven was thermally shielded by layers of tantalum foil and encased in a water-cooled jacket. Rough collimation of the Na beam was achieved by a 12mm-diameter aperture in the jacket. The Na beam was further collimated by a 6mm aperture ahead of the beam-crossing region (see fig. 2.2, item #3). The density of Na atoms in the beam-crossing region was $\sim 10^{13}$ particles/cm². This has been estimated using density equations for effusive flow which according to ref. [89] gives a good approximation for the supersonic expansion of alkali vapors. For an effusive molecular beam, the density of particles, ρ , decreases with distance, l , from the source according to $\rho = 0.25 \times \rho_0 \times (r/l)^2$, where ρ_0 is the density of particles directly ahead of the source and r is the radius of the orifice

The oven assembly was mounted on a translation stage in order that it could be moved away from near the pulsed-jet exit and the extraction electrodes of the TOFMS when measurements were not being made. This avoided the build up of large deposits of Na during the experiments which would block the expansion of the jet or disturb the homogeneity of the extraction field. The Na was deposited on a Cu block whilst the oven was in this position.

A pulsed valve driven by a solenoid (General Valve Corp.) with a nozzle diameter of 0.8mm, in a 1mm-length channel, was used to generate a supersonic beam of the halide in an inert-carrier gas. The supersonic expansion provided a vibrationally and rotationally-cold molecular beam. The backing pressure and the composition of the gas mixture which gave an optimal concentration of complexes was found to depend on the identity of the halide molecule. Typical backing pressures

used were between 250kPa (1900 torr) and 400kPa (3000 torr). The flow of the substrate and carrier gas into the gas line was controlled by precision needle valves. The partial pressure of gases were measured by a baratron pressure gauge (MKS Instruments, Inc. 222B) with digital readout (PDR-D-1). Once the gas line leading to the pulsed valve was filled with the appropriate gas mixture, complexes could be generated for a sufficiently long time before the mixture had to be replaced. However, it was found that the concentration of complexes decreased long before there was a drop in the backing pressure. This suggests that the halide/rare gas mixture separated in the gas line leading to the pulsed valve.

The pulsed valve was triggered by an IOTA ONE driver (General Valve Corp.) with pulse lengths between 250-300 μ s. It was found necessary to adjust the length of the trigger pulse after a large number of repetitions in order to compensate for changing characteristics of the pulsed-valve opening. This was caused by deformation of the poppet and the restoring springs inside the valve. At pulse lengths much longer than 300 μ s, the complex signal was significantly reduced due to the increase in gas pressure in the region of the beam and jet crossing point. The pulsed valve was typically operated at repetition rates between 5 and 20Hz.

The most important difference in the molecular-beam geometry of the current experiment, compared with that used in earlier experiments (see ref. 19), is the location of the crossing region. In the present arrangement, the distances from both nozzle exits to the geometric crossing has been significantly reduced. In the case of the pulsed valve, the reduction in the distance was from 6 mm to 0.5 mm and, for the

oven, it was from ~40 mm to 22 mm. In addition, the previous distance of ~70 mm from the crossing point to the ionization-laser beam was reduced to 15 mm in this work. A beam-crossing region close to both nozzle exits will give rise to a high local density of the alkali metal, halide and inert gas. This will result in more collision events in which complexes can be formed. The shorter distance from the beam-crossing region will also give rise to a higher density of complexes in the laser-interaction region.

The technique of using crossed-molecular beams for forming van der Waals complexes was pioneered by Hertel and co-workers in their studies of $\text{Na}\cdots\text{H}_2\text{O}$ [86]. Figure 2.3 illustrates a typical photoionization TOF mass spectrum, produced with the apparatus described here, in which a mixture of water and methanol in Ar-carrier gas was expanded through the pulsed valve. Thus formation of complexes $\text{Na}\cdots(\text{OH}_2)_n$ in the crossed-molecular beams, first identified by Hertel's group, has been reproduced in this laboratory.

In the present work, complexes of Na with methyl halide and hydrogen halide molecules were studied, i.e. $\text{Na}\cdots\text{XR}$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}, \text{R}=\text{CH}_3$; $\text{X}=\text{F}, \text{R}=\text{H}$). The efficiency of complex formation depended on the identity of the carrier gas, the composition of the gas mixture and the backing pressure of the pulsed valve. The optimal conditions were found to be different for each complex and they will be described individually in subsequent chapters. Experimental factors which affect the complex-signal intensity will be discussed in the following paragraphs and the system $\text{Na}\cdots\text{FH}$ will be used for illustration.

Figure 2.3: TOF mass spectrum obtained by expansion of a mixture of CH_3OH and H_2O in Ar-carrier gas through the pulsed valve. The complexes $\text{Na} \cdot \text{OH}_2$ were previously observed by Hertel and co-workers in a similar experimental apparatus[86].

Excitation laser, L_1

The van der Waals complexes formed in the crossed-molecular beams were carried by the momentum of the expanding jet a distance of 15mm into the laser-interaction region. The complexes were excited by a tunable-dye laser (L_1 ; Lambda Physik, LPD3000) pumped by an excimer laser (Lambda Physik, LPX200). The excimer-pump laser was operated with a Xe/HCl/Ne gas fill. The exciplex, (XeF)*, formed in the laser cavity gives rise to emission at 308nm. The path of L_1 is illustrated in fig. 2.4. The excitation laser, L_1 , was triggered after a delay of 400-800 μ s following the trigger to the pulsed valve. The laser entered chamber A through a series of light baffles in a side arm. This arrangement was designed to collimate L_1 and reduce the amount of scattered light in the laser-interaction region.

The pulse energy of L_1 was controlled by means of a prism attenuator (Newport Model 935-5). A beam splitter was used to reflect ~10% of the laser power of L_1 onto a Joule meter (Molelectron Model J3-02DW), the signal from which was amplified and measured by a home built sample-and-hold circuit.

Excitation of the vdW complexes will be shown to take place across a broad wavelength range, between 840 and 380nm. This region was scanned using a selection of laser dyes (Exciton) in organic solvents; LD 800 (DMSO, 860-780nm), Pyridin 2 (DMSO, 790-695nm), Oxazin 170 (Methanol, 728-660nm), Rhodamin 101 (Methanol, 672-614nm), Rhodamin B (Methanol, 644-588nm), Coumarin 153 (Methanol, 600-522nm), Coumarin 307 (Methanol, 553-479nm), Coumarin 102

Figure 2.4: Top view of the reaction chamber indicating the path of the excitation laser, L_1 , and the ionization laser, L_2 .

(Methanol, 510-460nm), Coumarin 120 (Methanol, 462-423nm), BisMSB (Dioxane, 428-414nm) and PBBO (Dioxane, 420-386nm).

Ionization laser, L₂

The vdW complexes which remained unreacted after laser L₁ was fired were subsequently ionized by an excimer laser, L₂ (Lumonics, TE 861 T-3). The arrangement is illustrated in fig 2.4 where the ionization laser, L₂, was counterpropagated to the excitation laser, L₁. The beam of L₂ was focused mildly by a f=1000 mm lens to fit its geometric cross section into an area within the diameter of the excitation laser, L₁. The duration of both laser pulses was ~20ns. The ionization laser L₂ was fired alone and then subsequent to the excitation laser, L₁, after a delay of ~250ns, to estimate the depletion intensity.

Laser L₂ was operated with a Kr/F₂/He gas fill. The exciplex, (KrF)*, formed in the laser cavity gives rise to emission at 248nm. The power of L₂ was measured, similarly to L₁, using a beam splitter to reflect a weak beam onto a separate Joule meter. The signal from which was amplified and measured by a second sample-and-hold circuit. The ionization laser, L₂, entered chamber A from the opposite side to L₁ through a side arm containing light baffles.

A typical TOF spectrum produced from the expansion of a mixture of ~2% HF in He carrier gas at 250kPa backing pressure is illustrated in figure 2.5. The broken line indicates the ion signal when the ionization laser, L₂, was fired alone. A strong signal from Na₂ and a weak signal from Na is evident to either side of the Na··FH peak. The sodium dimer, Na₂ (I°= 4.89eV[93]), can be ionized by a single photon at 248nm, but Na

Figure 2.5: TOF mass spectrum obtained by expansion of a 2.5% mixture of HF in He at 250kPa backing pressure through the pulsed valve. The dotted line gives the ion signal obtained when L_2 ($\lambda_2=248\text{nm}$) was fired alone and the solid line gives the ion signal when L_1 ($\lambda_1=660\text{nm}$) was fired prior to L_2 .

($I^\circ = 5.14\text{eV}$ [93]) must be ionized via a two-photon process. The Na peak has a broad companion peak to longer flight times which is likely to have arose from fragmentation of $[\text{Na}\cdots\text{FH}]^+$ or Na_2^+ along the ion-flight path. In addition, the presence of H_2O as an impurity in the gas inlet gave rise to a weak signal which is identified as $\text{Na}\cdots\text{OH}_2$ complexes in the TOF spectrum.

The solid-line in the TOF spectra in fig. 2.5 indicates the extent of depletion of the $\text{Na}\cdots\text{FH}$ complexes following excitation by laser, L_1 , which was set at a wavelength of $\lambda_1 = 660\text{nm}$ and fired prior to the ionization laser, L_2 . The wavelength of L_1 was chosen to lie within a broad photodepletion region for this complex (550-830nm -see chapter 4); the chosen λ_1 in fig. 2.5 is merely illustrative. The Na and Na_2 signal were unaffected by the excitation laser, L_1 .

Ion collisions that form species $n-1$ from a complex n appear as a broad tail at longer flight times attached to the signal of complex $n-1$. This effect was discriminated against when measuring photodepletion by the gated integration of the complex signal within a narrow time window. The measured photodepletion is not affected by collision-induced dissociation, since this process is operative with and without the depletion laser, L_1 .

The ionization energy for $\text{Na}\cdots\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) and $\text{Na}\cdots\text{FH}$ complexes are believed to lie below 5.0eV (the energy corresponding to the wavelength, $\lambda_2=248\text{nm}$, of L_2). An ionization energy of 4.3eV was determined by Wang et al[19] for $\text{Na}\cdots\text{FCH}_3$. The dependence of the ion signal on the ionization-laser fluence, F_2 , was measured for each

complex. A linear relationship was observed in each case, consistent with a single-photon process, and is illustrated in fig. 2.6 as a log plot for the system $\text{Na}\cdots\text{FH}$.

The logarithm of the photodepletion signal, $-\ln(N/N_0)$, is shown as a function of the excitation-laser fluence, F_1 , in fig. 2.7. A linear increase up to 4 mJ/cm^2 was in good agreement with a previous measurement for $\text{Na}\cdots\text{FCH}_3$ [19]. Saturation was observed towards higher laser fluence. In these experiments, the excitation-laser fluence was stabilized in the range $2.5\text{-}3.0 \text{ mJ/cm}^2$ when recording the depletion of the various complexes as a function of the excitation wavelength. Consequently, the measured depletion was much less than that used for illustration in fig. 2.5.

Time-of-flight mass spectrometer

An acceleration field of 25 Vcm^{-1} was used to extract the ions from chamber A into the time-of-flight mass spectrometer (TOFMS). This first acceleration stage of the TOFMS was composed of two parallel stainless-steel plates which were separated by 1.5 cm . A positive potential was applied to the first plate (item #1 in fig 2.1) whilst the second plate was earthed (item #2 in fig. 2.1). The ions formed by L_2 traveled over a distance of about 3 mm to pass through a $3\text{ mm}\times 10\text{ mm}$ slot in a box-shaped extension of the second electrode. The homogeneity of the electric field used to extract the positive ions was improved by attaching a metal plate with teflon isolation to the bottom face of the pulsed valve. A stainless-steel wire was then placed below this plate and the laser-interaction region, running parallel with the extraction electrodes. The

Figure 2.6: Power dependence of the ionization signal for Na·FH complexes measured at 248nm. The straight line is a linear least-squares fit to the data points. The slope of the line is 1.02.

Figure 2.7: Power dependence of the depletion signal for the complex Na·FH measured at $\lambda_1 = 660$ nm. A least-squares fit to the data is given for the linear region below 2 mJ/cm².

pulsed-valve plate was supplied with half the potential on the repeller plate and the wire was given a potential which was tunable externally between the plate and the repeller voltage. The wire worked as a deflector, permitting adjustment of the perpendicular-velocity component of the ion beam, so as to obtain optimum alignment along the TOF axis. This arrangement is not illustrated in fig. 2.1.

The extraction field was pulsed with a time delay of approximately 250ns from the ionization laser, L_2 . Spurious ion signals were observed when a constant extraction field was applied to the electrodes. These signals likely arose from electron-impact ionization of atoms and molecules by electrons produced from a gas discharge triggered by photoionization, or alternatively by photoemission at sodium covered surfaces (the final energy of the electron, accelerated by the electric field, would be $\sim 25\text{eV}$). The complex-ion intensity as a function of the time delay between laser L_2 and the extraction field is shown in fig. 2.8 for $\text{Na}\cdots\text{FH}$ complexes. The ion intensity remained constant between delay times of 0 and 600ns. Longer time delays showed an intensity decrease because ions were drifting away from the aperture of the second electrode and, therefore, the acceptance region of the TOFMS.

The second acceleration stage of the TOFMS was located in chamber B. The ions traveled along a short flight tube (item #3 in fig. 2.1) and were then further accelerated to a kinetic energy of $\sim 2200\text{eV}$ and focused simultaneously by a series of twelve equally-spaced electrodes with apertures of decreasing diameter (item #4 in fig. 2.1). Trajectory simulations for the ion path, employing the MacSimion computer program[94], assisted in selecting values for the electrode voltages in

Figure 2.8: The ion-signal intensity for $\text{Na}^+\cdot\text{FH}$ complexes as a function of the time delay between the ionization laser, L_2 and the onset of the pulsed-extraction field.

chamber B. For instance, it was found that a linear increase in the magnitude of the potential on the electrodes at the early stages of the flight path in chamber B would provide a well-collimated ion beam and a large potential on the penultimate electrode would then provide a focused ion beam on the detector. The magnitude of the electrode voltages were adjusted externally (Spellman RHSR 5PN50 power supply with a home-built potential divider) in order to optimize the complex-signal intensity. Typical potentials for the acceleration electrodes are indicated in table 2.1.

The 60cm-flight tube in chamber C (item #5 in fig 2.1) was kept at a negative potential equal to that for the final acceleration electrode in order to maintain a focused ion beam. The flight tube was terminated by a wire mesh. The ions were detected by a dual set of microchannel plates (MCP, Galileo, 1396-0060 matched pair) in a chevron configuration, mounted in a shielded housing. This arrangement is illustrated in fig. 2.1 (item #7), in which the ions first have to pass through a grounded wire mesh which protects the MCP from intense signals. A potential of 1000V was supplied across each microchannel plate (Bertan power supply, model no. 215). The electrons generated by the MCP were measured at the final electrode in fig. 2.1. The ion-signal was preamplified by a factor of 10 using a 300 MHz amplifier (Philips Scientific Model 6950).

Time-of-flight spectra, consisting of 2048 data points taken at intervals of 40ns, were recorded on a 100 MHz oscilloscope (Tektronix 2232). Averaging of the ion signal from spectra recorded in sequence (using at least 100 laser shots) provided a good signal-to-noise ratio. The final spectrum was collected by a microcomputer (Macintosh SE/30). For

Table 2.1: Electrode voltages for the TOFMS

Electrode	Potential /V	Electrode	Potential /V
1	+25	4 (f)	-1400
2	0	(g)	-1600
3	-200	(h)	-1800
4(a)	-400	(i)	-2000
(b)	-600	(j)	-2200
(c)	-800	(k)	-4000
(d)	-1000	(l)	-2200
(e)	-1200	5	-2200

intensity measurements of the ion signal for a selected vdW complex, a boxcar gated integrator and signal averager (Stanford Research SR250) was used. The analogue signal was then converted to a digital signal and the data was collected by the microcomputer via an interface (SR245). The same interface was also used as a means to collect measurements of the excitation and ionization-laser fluence, F_1 and F_2 , from the sample-and-hold circuits.

A "mass gate" (item #6 in fig. 2.1) was inserted directly after the flight tube in order to protect the detector assembly from saturation caused by a strong Na_2 signal when ion signals corresponding to masses above 46 a.m.u. were being measured. The gate consisted of a wire mesh which was pulsed from the flight-tube potential to a sufficiently high voltage (Spellman RHSR 5PN50 power supply/ Directed Energy Inc. GRX-E high-voltage pulser) and at a delay from the ionization laser appropriate to reflect unwanted ions of low mass.

Timing circuitry

The trigger signal to the pulsed-valve driver, excitation laser (L_1), ionization laser (L_2) and mass gate was provided by a 5-channel digital-delay/ pulse generator (Stanford Research, DG535). In addition, the output from the DG535 directly supplied the extraction electrodes with a voltage pulse which could be varied from 0 to +40V (typically a value of ~+25V was used- see table 2.1). A series of measurements were performed to confirm that the depletion signal did not reflect the time delay between successive operations.

The optimal time delay between the pulsed valve and the laser interaction was found to be dependent on the backing pressure for the pulsed valve. The dependency is illustrated in fig 2.9 for Na $\cdot$ FH complexes formed using a partial pressure of $\sim$ 5kPa for HF. The ion signal represents the complex-signal intensity when the ionization laser, L₂, was fired alone. Initially, the signal increased with the backing pressure of the pulsed valve. A maximum signal intensity was reached at an intermediate pressure, which for the example of Na $\cdot$ FH complexes was at $\sim$ 170kPa using a 650 μ s time delay between the pulsed valve and the ionization laser. A further increase of the backing pressure then resulted in a decrease of the complex-signal intensity. However, this decrease could be partially offset by either reducing or increasing the time delay between the pulsed valve and the ionization laser. The decrease in the ion signal at high backing pressures arises from collisional dissociation of the complexes in the extraction region of the TOFMS due to the high density of particles in the gas cloud formed from the expansion of the pulsed-supersonic jet. At shorter or longer delay times, observation is made of the early or late edges of the gas pulse, in which the density of particles is lower.

In these experiments, the time delay between the pulsed valve and the laser interaction was optimized in the region between 400-800 μ s according to the backing pressure used. The optimal ion-signal for different Na $\cdot$ XR complexes was found to require different pairs of values for the backing pressure and the time delay between the pulsed valve and the ionization-laser pulse. For example, Na $\cdot$ XCH₃ (X=F, Cl, Br) complexes were generated by the expansion of a concentrated mixture of

Figure 2.9: The ion-signal intensity for the complex $\text{Na}\cdot\text{FH}$ as a function of the backing pressure for the pulsed valve and the time delay between the ionization laser, L_2 , and the pulsed valve.

XCH₃ in Ar carrier gas, at high backing pressure and using a relatively long time delay. However, the Na[•]·FH complex was generated by the expansion of a dilute mixture of HF in He carrier gas at a lower backing pressure and a shorter time delay. The individual conditions used to generate each complex will be described in the subsequent chapters.

The depletion signal N/N_0 is shown in fig. 2.10 as a function of the time delay between the excitation laser, L_1 , and the ionization laser, L_2 for Na[•]·FH complexes. Depletion of the ion signal was not observed if the excitation laser, L_1 , was fired after the ionization laser L_2 . This indicates that photoinduced reaction of the ionic complex, Na⁺·FH, does not occur in this region of excitation wavelengths. The depletion of neutral complexes were found to be independent of the time delay across a region of 100-400ns (see fig. 2.10). The definition of time delay $\Delta t = 0$ is broadened by a time jitter between both lasers (~50 ns), as well as the duration of the laser pulses (~20 ns). At times in excess of 400ns, the measured value of N/N_0 increases due to replenishment of the Na[•]·FH complex in the extraction region of the TOFMS. A time-delay of 250ns was used between L_1 and L_2 for these experiments.

2.2 Data Analysis

Mass spectra

The mass number of the ions which correspond to a particular peak in the TOF spectrum were identified according to the flight time. Since the kinetic energy, $E_{tr} = mv^2/2$, of each accelerated ion will be the same, the ion mass will be related to the flight time, t , by

Figure 2.10: Normalised ion-signal intensity for the complex $\text{Na} \cdots \text{FH}$ as a function of the time delay between the ionization laser, L_2 , and the excitation laser, L_1 .

$$\sqrt{m} = c \times (t - t_0) \quad (2.1)$$

where t_0 is defined by the time at which the extraction field is pulsed and c is a constant to be determined using known mass numbers and their flight times. The plots of $\sqrt{m}$ versus the flight time could be fitted into the form of eq. 2.1, which in turn converted the TOF spectrum into a mass spectrum.

Photodepletion measurements

The dynamics of the excited-state reaction were investigated by observing the depletion cross section for the complex as a function of the excitation wavelength. The concentration of complexes before and after irradiation with the excitation laser was measured, and the depletion cross-section σ_{dep} in $\text{\AA}^2$ was calculated using the equation,

$$\sigma_{\text{dep}} = (1/F_1) \ln (N_0/N) \quad (2.2)$$

where N_0 and N represent the ion signal of the vdW complex before and after photodepletion, and F refers to the excitation-laser fluence in photons/ $\text{\AA}^2$. The depletion cross section provided a normalised scale which could be used to compare the extent of depletion for different complexes and also as a function of the excitation wavelength. The origin of eq. 2.2 will be discussed in the following paragraphs.

The depletion of the complex $\text{Na}\cdot\text{XR}$ following excitation with a wavelength, λ_1 (hence ν_1), is given by

If it is assumed that all the excited complexes react, leading to depletion of $\text{Na}\cdots\text{XR}$ then

$$\frac{dN}{dt} = - \sigma_{\text{dep}} I_1 N \quad (2.4)$$

where N is the concentration of complexes (or the ion signal in the TOF spectrum), I_1 is photon-flux of the excitation laser (L_1) in photons/(cm^2s) and t is the time elapsed during the L_1 -laser pulse (i.e. $0 < t < T$ where T is the period of the laser pulse). After L_1 has fired, the concentration of complexes will have been reduced from N_0 to N , and an explicit expression for the depletion cross section in eq. 2.2 can be calculated using

$$\sigma_{\text{dep}} = - \int_{N_0}^N \frac{dN}{N} / \int_0^T I_1 dt \quad (2.5)$$

$$\text{in which } \int_{N_0}^N dN = \ln(N_0/N), \text{ and } \int_0^T I_1 dt = F_1$$

where F_1 is the fluence of L_1 in mJ/cm^2 . The applicability of eq. 2.2 to the systems studied here is confirmed by the linear plot in fig. 2.7 for a photon fluence of L_1 below $4 \text{ mJ}/\text{cm}^2$. This equation bears similarity to the Beer-Lambert law. In the latter example, the intensity of the incident radiation, I , as it passes a distance x through a sample of concentration, N , is attenuated according to

$$\frac{dI}{dx} = - \kappa I_1 N \quad (2.6)$$

Equations 2.4 and 2.5 can be shown to be equivalent. The derivative dI/dx will be proportional to dN/dx and dx/dt is a constant (the speed of light). It follows that dI/dx will be proportional to dN/dt . The

proportionality constant, κ , in eq. 2.6 is then replaced by the depletion cross section, σ_{dep} , in eq. 2.4. Alternatively, eq. 2.6 can be solved directly to give the familiar expression,

$$\log_{10} (I_0/I) = \epsilon N l$$

where ϵ is the molar absorption coefficient and l is the path length through the sample of concentration, N .

The depletion cross section of the complexes measured in the TOF mass spectra were found to fluctuate between measurements by $\pm 5\%$. This is likely to arise from different concentrations of the complexes being formed in the molecular beam due to an instability in the expansion of the halide molecules from the pulsed-valve source. The measurements of σ_{dep} were typically averaged over 100 repetitions of the L_1 and L_2 laser pulses. In addition, from day to day, the magnitude of the cross section in the photodepletion spectrum (measured as a function of the excitation wavelength) was found to vary by $\pm 20\%$. The origin of this uncertainty in the measurements was likely to have been due to slight changes in either the alignment of the overlapping laser beam or the homogeneity of the excitation-laser beam.

The measurement of the depletion cross-section for a complex as a function of the wavelength of the excitation laser will be referred to as an action spectrum. The spectra obtained will be described for $\text{Na}\cdot\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) in chapter 3 and for $\text{Na}\cdot\text{FH}$ in chapter 4. The action spectra will be shown to provide information on the dynamics for the excited-state reactions, $\text{Na}^* + \text{XR} \rightarrow \text{NaX} + \text{R}$.

2.3 Summary

The complexes $\text{Na}\cdots\text{XR}$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$, with $\text{R}=\text{CH}_3$, $\text{X}=\text{F}$, with $\text{R}=\text{H}$) were formed using a crossed-beam apparatus in a vacuum chamber. The Na atomic beam was generated from an oven source and the halide molecular beam from a pulsed-supersonic expansion. The complexes were identified by photoionization time-of-flight mass spectrometry. This section described experimental methods used to measure the depletion of the complexes as a function of the excitation wavelength, λ_1 . The dependence of the ion signal and the depletion on experimental factors such as the excitation and ionization-laser fluence, the time delay between successive operations and the expansion characteristics of the pulsed valve have been described. Conversion of the depletion to a cross-section, σ_{dep} , permits the comparison of the measurements for different complexes and different excitation wavelengths, λ_1 .

Chapter 3 Photodepletion of $\text{Na}\cdots\text{XCH}_3$; $\text{X}=\text{F}, \text{Cl}, \text{Br}$

This chapter presents a comparative study of the photoinduced charge-transfer dissociation reaction in $\text{Na}\cdots\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) van der Waals complexes. The depletion cross-section for these complexes was recorded as a function of the excitation wavelength across a wide range from 750 to 390nm. The resulting action spectra were found to consist of a series of broad peaks for each system.

Photoexcitation of $\text{Na}\cdots\text{XCH}_3$ accesses a potential-energy surface (PES) which would be traversed by the reaction of an excited state of Na with the methyl-halide molecule, i.e. $\text{Na}^* + \text{XCH}_3 \rightarrow \text{NaX} + \text{CH}_3$. Monochromatic excitation forms the transition-state (TS), $[\text{Na}^*\cdots\text{XCH}_3]^\ddagger$, in a selected-starting configuration. The photodepletion peaks have been assigned according to the atomic state of Na^* which correlates with the TS formed in each wavelength region. The assignment was assisted by *ab initio* calculations, performed by others in this laboratory (see ref. 96), of the potential energies for the excited-state complexes.

The depletion cross-section is a product of two processes as there is, first, photoexcitation of the complex to an electronically-excited state, $\text{Na}\cdots\text{XCH}_3 + h\nu_1 \rightarrow [\text{Na}^*\cdots\text{XCH}_3]^\ddagger$, and, subsequently, a reaction which leads to removal of the complex, $[\text{Na}^*\cdots\text{XCH}_3]^\ddagger \rightarrow \text{Products}$. The cross-section for each photodepletion peak integrated across the appropriate energy interval will be compared to the oscillator strength, obtained by *ab initio* calculation[96], for the appropriate electronic transition in the complex. This will indicate the probability with which depletion occurs on each excited-state PES.

The measured photodepletion peaks were found to contain reproducible structure which was believed to arise from vibrational levels of the complex. *Ab initio* calculation of the structures of the ground and electronically-excited states, along with the frequencies for the vibrational modes[96] will be used to assist assignment of the observed progressions.

An earlier publication from this group focused on a study of the series of complexes $\text{Na}\cdots(\text{FCH}_3)_n$, $n=1$ to 4, which were formed using a different geometry for the crossed-molecular beams[19]. At this time, complexes with BrCH_3 were not identified and the concentration of complexes with ClCH_3 was found to be insufficient to yield reliable photodepletion measurements. The improved efficiency for complex formation with the current geometry (see chapter 2) has enabled a study of all three methyl-halide systems. An improved signal-to-noise ratio has identified reproducible structure within the photodepletion peaks for $\text{Na}\cdots\text{FCH}_3$ which was not sufficiently resolved in the previously published spectrum[19]. In addition, the assignment of the photodepletion peaks previously proposed for the system $\text{Na}\cdots\text{FCH}_3$ has been revised based on more accurate *ab initio* calculations[96].

3.1 Identification of complexes

The van der Waals (vdW) complexes $\text{Na}\cdots\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) were formed in the crossed-molecular beams using an expansion of XCH_3 in an inert carrier gas from the pulsed-jet source, as described in section 2.1. A total backing pressure for the pulsed valve between 300 and 400kPa was found to give a strong complex signal.

Although the gas mixture introduced into the inlet of the pulsed valve contained a high concentration of XCH_3 molecules (30-70%), it was very likely that considerable separation of this mixture occurred in the gas line prior to the expansion. This was in accord with the observation that the complex-signal intensity decreased with time although the backing pressure of the pulsed valve still remained high (see section 2.1). The composition of the mixture expanded from the pulsed valve was expected to have contained a concentration of the methyl halide which was much less than that introduced into the gas line initially.

For FCH_3 expansion, the choice of the inert gas was not important for complex formation (He or Ar being used), but for $ClCH_3$ and $BrCH_3$ double the intensity of complex was obtained using argon. The inert gas not only played a role in cooling the halide molecules to low rotational and vibrational temperatures, but also acted as a third body to carry away the excess collision energy in order to stabilize the vdW complex.

There may be a number of reasons for the improved complex intensity using argon, instead of helium, as the carrier gas. A likely mechanism for formation of $Na \cdots XCH_3$ would involve displacement of the rare-gas atom, $Rg = He$ or Ar , from the weakly bound $Rg \cdots XCH_3$ cluster by a colliding Na atom. The precursor complex, $Rg \cdots XCH_3$, is expected to be present in the expansion region of the supersonic jet. The existence of these clusters would be enhanced at the high backing pressure used in the pulsed-valve source. The departing Rg atom will carry away the excess collision energy as the complex $Na \cdots XCH_3$ is formed. An argon atom has a larger polarizability than a helium atom and, therefore, the

van der Waals interaction with a halide molecule will be stronger, giving a higher concentration of the precursor complexes.

In addition, a supersonic expansion in Ar carrier gas will give rise to a lower collision energy for $\text{Rg}\cdots\text{XCH}_3$ with Na atoms than an expansion in He carrier gas. The velocity of a supersonic expansion of XCH_3 in excess carrier gas would be $\sim 500\text{m/s}$ in Ar and $\sim 1700\text{m/s}$ in He [92]. This gives rise to a perpendicular collision energy of $\sim 0.2\text{eV}$ for $\text{Na} + \text{Ar}\cdots\text{XCH}_3$, compared with $\sim 0.4\text{eV}$ for $\text{Na} + \text{He}\cdots\text{XCH}_3$. It appears that these factors are important in forming the van der Waals complexes $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ which are more weakly bound than $\text{Na}\cdots\text{FCH}_3$ (see below).

Photoionization time-of-flight (TOF) mass spectra for the systems $\text{Na} + \text{XCH}_3$, $\text{X} = \text{F}, \text{Cl}, \text{Br}$ were recorded using identical mixtures for the halide in Ar-carrier gas at a backing pressure of 300kPa . In addition, an expansion mixture containing a much lower concentration of the halide with Ar was used for ICH_3 to record a TOF spectrum for the system $\text{Na} + \text{ICH}_3$. The composition of this mixture was limited by the low vapor pressure of ICH_3 which is a liquid at room temperature. For this substrate, the carrier gas was bubbled through a glass reservoir containing ICH_3 , as described in ref. 95. These TOF spectra, recorded under otherwise identical conditions, are represented by the dotted lines in fig. 3.1. The mass spectra show ion signals for the complexes $\text{Na}\cdots(\text{XCH}_3)_n$ with n ranging for $\text{X} = \text{F}$ from $n = 1-5$, for $\text{X} = \text{Cl}$ from $n = 1-4$, and for $\text{X} = \text{Br}$ from $n = 1-2$. However, only a very small ion signal was recorded for the $\text{Na}\cdots\text{ICH}_3$ complex. Instead, the mass spectrum for the $\text{Na} + \text{ICH}_3$ system shows the reaction products, NaI and Na_2I .

Figure 3.1: TOF ion spectra for the systems $\text{Na}+\text{XCH}_3$, $\text{X}=\text{F}, \text{Cl}, \text{Br}, \text{I}$. All peaks are labeled by the corresponding uncharged species. The broken line represents the spectra obtained when L_2 (248nm) was used alone. The solid line gives the depleted signal intensities when L_1 (666nm, $\sim 8\text{mJ}/\text{cm}^2$) was used shortly before L_2 to excite the complexes.

The complex-signal intensity reflects the photoionization efficiency and the probability for complex formation. The yield of complexes for the case of $n=1$ was found to be in the ratio $X=F:Cl:Br$ of 10 : 1 : 1. This should be compared with the relative yield 160:1 for $X= F: Cl$ observed with the previous experimental set-up[19]. The current molecular-beam geometry appears to enhance the efficiency for formation of more weakly bound complexes.

Water impurities in the gas line leading to the pulsed valve led to the formation of the $Na\cdots OH_2$ complexes which were identified by Hertel and co-workers in their crossed-beam experiment[86], but complexes of Na_2 with $ClCH_3$ were not detected. This latter result is in contradiction to previous experiments performed in this laboratory[17] which found a sufficient intensity of these sodium-dimer complexes to study their photodepletion. The intensity of the single sodium complex, $Na\cdots ClCH_3$ in the previous work was, by contrast, too low to yield reliable photodepletion results.

The absence of $Na\cdots ICH_3$ complexes was likely to be due to reaction occurring in the crossing region of the molecular beams in the absence of photoexcitation. The mean collision energy for the $Na+CH_3I/Ar$ system was ~ 0.2 eV. In a crossed-beam experiment for $K+ICH_3$ performed by Bernstein and coworkers[44] a translational energy barrier of 0.039 eV was measured for reaction. The equivalent barrier for reaction with Na is expected to be only slightly higher, but ~ 0.2 eV, since the energy barrier for reaction in the electronic-ground state is likely to be more dependent on the identity of the methyl-halide molecule than on the alkali-metal atom[97]. The origin of the barrier has been described as arising from the

energy required to stretch the X-CH₃ bond in order to obtain a positive electron affinity such that charge-transfer to give ⁻X-CH₃ can then occur. Using this procedure a simple calculation using empirical potential-energy curves for CH₃I and CH₃I⁻ gave a value of 0.026eV for this energy[48].

The solid line in fig. 3.1 shows the extent of photodepletion for the different systems Na[·]XCH₃ following excitation by laser L₁, which is fired ahead of the ionization laser, L₂, at a wavelength, λ₁, of 666nm. The wavelength of L₁ was chosen to lie in the broad photodepletion region identified in earlier experiments[19] and its value is merely illustrative. The laser fluence for L₁ and L₂ were kept constant for all measurements. For the purpose of fig. 3.1, the power of L₁ was high (~8mJ/cm²) resulting in depletion of ~50%. However, in the photodepletion studies the power of L₁ was kept low enough to give a depletion intensity <10% in order that the Beer-Lambert law would be applicable (see section 2.2). The depletion of the complex was largest at 666 nm for X=Cl. Photodepletion was also observed for the complexes Na[·]·(XCH₃) with n>1, as well as for Na[·]·OH₂.

The mass spectra in fig. 3.1 show the reaction product, NaI, from Na+ICH₃ but not from photoinduced reactions with the other methyl halide molecules. However, we would not expect to observe NaX (X= F, Cl, Br) by laser ionization at 248nm. With ionization potentials for sodium halide molecules of approx. 8.5 eV for NaBr, 8.9eV for NaCl and 9.5 eV for NaF[98], at least two photons of 248 nm (5 eV) would be required for photoionization. At typical ionization-laser fluences, F₂, of 30 mJ/cm² a two-photon process would be unlikely to be observable in

the absence of a resonant intermediate state. The low electronically-excited states of alkali-halide molecules are, in general, believed to be dissociative[99]. However, the potential energy curves for excited states of NaI obtained by Davidovits et al.[100] show one state with a shallow minimum of ~ 0.2 eV, which is accessible with a 5 eV photon from certain excited vibrational levels ($v=4-8$) of the ground state. We believe that we have populated these vibrational levels by way of the exothermic reaction $\text{Na} + \text{ICH}_3 \rightarrow \text{NaI} + \text{CH}_3$, for which, $\Delta H^\circ = -17.0$ kcal/mol ($D^0(\text{I-CH}_3) = 57$ kcal/mol[101] and $D^0(\text{NaI}) = 72.7$ kcal/mol[93]). The dependence of the signal on the ionization-laser power is shown in fig. 3.2. A dependence much greater than unity indicates a multiphoton process.

The small Na^+ signal in fig. 3.1(d) showed the same intensity dependence of 1.7 on the laser fluence as did NaI^+ (see fig. 3.2). The Na^+ was thought to arise from single-photon excitation of NaI ($v > 8$) to yield $\text{Na}^*(2P) + \text{I}(2P_{3/2})$, followed by photoionization of $\text{Na}^*(2P)$ by a second photon. The Na_2I^+ peak in fig. 3.1(d) exhibited a power dependence which suggests that a single-photon ionization process is likely to have occurred (see fig. 3.2). It is likely to be due to direct ionization of Na_2I produced in the reaction of Na_2 with ICH_3 . The ionization potential of Na_2I is expected to lie below that for Na_2F which is 3.5 ± 0.15 eV [102]. Hence a single photon of 248nm (corresponding to 5.0eV) will suffice to give Na_2I^+ , as observed.

In the case of the $\text{Na} \cdots \text{BrCH}_3$ complexes we found a small intensity of Na^+ due to the combined presence of the excitation ($\lambda_1=666\text{nm}$) and ionization ($\lambda_2=248\text{nm}$) laser radiation. The excitation energy ($E_1=1.9\text{eV}$) in this instance is below that required to access the lowest

Figure 3.2: Power dependence of the ion signals for Na, NaI and Na₂I in the TOF spectra. The ionization wavelength was 248nm. The straight line represents a linear least-squares fit to the data points.

electronically-excited state of sodium (i.e. the D-line excitation at 2.1 eV) which implies that the excited complex cannot fragment into an excited-sodium atom, $\text{Na}^*(3p^2P)$. This rules out ionization of excited-state Na^* , which is formed by L_1 , as the origin of the Na^+ signal. A more likely explanation is ionization of the excited-state complex, leading to a high vibrational state of the complex ion $(\text{Na}^+\cdots\text{BrCH}_3)^+$ which subsequently predissociates to give $\text{Na}^+ + \text{BrCH}_3$. The lifetime of the excited-state complex must exceed 10^{-7}s in order that this two-photon process can occur.

We believe that the major route for photodepletion involves formation of the harpooning-reaction products, $\text{NaX} + \text{CH}_3$. The failed reaction pathway, leading to $\text{Na} + \text{XCH}_3$, is believed to be a minor route. The TOF spectrum for $\text{Na}\cdots(\text{XCH}_3)_n$ complexes in fig. 3.1 shows a strong signal for Na with two attached XCH_3 molecules. Depletion of the $n=2$ complex was also observed in the presence of the excitation laser. The photoformation of $\text{Na}\cdots\text{XCH}_3$ from van der Waals bond cleavage of $\text{Na}^*\cdots(\text{XCH}_3)_2$ was shown in earlier work not to be significant[19]. In which case, it seemed likely that the major channel for depletion of $\text{Na}^*\cdots\text{XCH}_3$ would also not involve van der Waals bond cleavage, in this case leading to $\text{Na} + \text{XCH}_3$. The two pieces of experimental evidence which were used to indicate that van der Waals bond breaking were not major pathways were;

(1) A linear plot of σ_{dep} for $\text{Na}\cdots\text{XCH}_3$ against the excitation-laser fluence, F_1 would not be obtained if depletion and formation of $\text{Na}\cdots\text{XCH}_3$, from respectively $\text{Na}^*\cdots\text{XCH}_3$ and $\text{Na}^*\cdots(\text{XCH}_3)_2$, were occurring

simultaneously. The power dependence of the depletion signal was shown to be linear for $\text{Na}\cdots\text{FCH}_3$ in ref. 19 (and for $\text{Na}\cdots\text{FH}$ in fig.2.7).

(2) It was possible to alter the ratio of $\text{Na}\cdots(\text{FCH}_3)_n$ for $n=2$ and $n=1$ which were formed in the crossed-molecular beams by changing the expansion conditions of the supersonic jet (such as the backing pressure or the composition of the gas mixture). Using this technique, it was found in ref. 19 that the photodepletion measurements for $\text{Na}\cdots\text{FCH}_3$ were invariant on the relative proportion of $n=1$ and $n=2$ complexes in the laser-interaction region. This should not be the case if the complex $\text{Na}\cdots\text{FCH}_3$ is being formed from photolysis of $\text{Na}\cdots(\text{FCH}_3)_2$.

Since the photoinduced reaction of the complexes appears not to involve van der Waals bond cleavage, the major channel for reaction of $\text{Na}^*\cdots(\text{XCH}_3)_n$ must be the pathway leading to the harpooning reaction products, $\text{NaX}\cdots(\text{XCH}_3)_{n-1} + \text{CH}_3$.

3.2 Structure of complexes

The ground-state geometries for $\text{Na}\cdots\text{FCH}_3$, $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ were fully optimized at the second-order Moller-Plesset (MP2) level by using Gaussian 92. A 6-31G** basis set was used for H, C, F and Cl atoms and an effective-core potential for Br. The standard basis set 6-31G* was used for Na, incorporating diffuse s, p and d wavefunctions. The details of this calculation, performed by others in this group, is described in ref. 96.

The minimum-energy structures were calculated to arise from attachment of Na at the X end of the halide molecule in accord with the

results of Trenery et al.[42]. This is also consistent with experimental observation of a large difference in ion yield, and therefore stability of the complex down the series $\text{Na}\cdots\text{FCH}_3$, $\text{Na}\cdots\text{ClCH}_3$, $\text{Na}\cdots\text{BrCH}_3$, namely, 10:1:1. This is readily understandable if the Na binds to the halogen, but not if Na binds to the organic radical which is the same for all three complexes.

The ground-state structure for the three complexes, $\text{Na}\cdots\text{XCH}_3$, ($\text{X}=\text{F}$, Cl and Br) determined by *ab initio* calculation are shown in fig. 3.3. It was found that each complex had a bent configuration, in which the bond angle decreased in the order $\text{Na}\cdots\text{F}-\text{CH}_3 > \text{Na}\cdots\text{Cl}-\text{CH}_3 > \text{Na}\cdots\text{Br}-\text{CH}_3$. There are two contributing factors which could give rise to the non-linear structure;

(1)The van der Waals bond could be considered to be formed from a donor-acceptor interaction in which the lone pair of electrons on X are donated to an empty 3p orbital on Na. The orientation of the electron pairs on X, which now consist of two bonding orbitals and two lone pair orbitals, would have distorted tetrahedral symmetry (such as in the H_2O molecule).

(2)The bonding in the ground-state complex will possess a significant degree of ionic character. Charge-transfer from Na into the antibonding orbital of XCH_3 will accumulate positive charge on Na and negative charge on C (the electron distribution in the antibonding orbital is the inverse of the bonding orbital). Long-range coulombic attraction between Na and C will provide an attractive contribution to bending in $\text{Na}\cdots\text{XCH}_3$.

Figure 3.3: Geometry and dissociation energy of the most stable configurations for the complexes $\text{Na}\cdots\text{XCH}_3$, $\text{X}=\text{F}, \text{Cl}, \text{Br}, \text{I}$, calculated using Gaussian92 at the CIS level[96]. The $\text{Na}\cdots\text{X}$ bond length in the complex can be compared with the equilibrium bond length in the alkali-halide molecule.

The complex $\text{Na}\cdots\text{FCH}_3$ has a much higher dissociation energy than $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$. The van der Waals bond lengths, $r(\text{Na-X})$, are compared with equilibrium bond lengths, $r_0(\text{Na-X})$, in the product alkali-halide molecules. The dissociation energy for the three complexes is reflected by the extension of the Na-X bond relative to the value in the equilibrium NaX molecule, i.e. $\Delta r(\text{Na-X})/r_0(\text{Na-X})$ where $\Delta r(\text{Na-X})=r(\text{Na-X})-r_0(\text{Na-X})$. This ratio increases (as the bond energy decreases) in the order $\text{X}=\text{F} > \text{Cl} > \text{Br}$. The van der Waals bond length in $\text{Na}\cdots\text{FCH}_3$ is particularly short, only 20% longer than the equilibrium bond length in the NaF molecule. However, the F-C bond in the $\text{Na}\cdots\text{FCH}_3$ complex, was stretched only slightly compared with the bond length in the FCH_3 molecule (in which $r_0(\text{F-C})$ is 1.382 Å[93]). For $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ the van der Waals bond lengths are much longer relative to the equilibrium bond lengths in NaCl and NaBr, and there is almost no stretch in the C-X bond length. The H-C-H angle and C-H bond length for the three complexes were determined to be essentially unaffected by the presence of the Na atom.

3.3 Photodepletion measurements

Photodepletion of the vdW complexes $\text{Na}\cdots\text{XCH}_3$, $\text{X}=\text{F}, \text{Cl}, \text{Br}$, was measured to either side of the Na D-line across an approx. $\sim 1.5\text{eV}$ range of excitation energies. The action spectra are the depletion cross-section as a function of the excitation wavelength, obtained by tuning L_1 at a fixed ionization wavelength for L_2 .

The action spectra for $\text{Na}\cdots\text{XCH}_3$, $\text{X}=\text{F}, \text{Cl}, \text{Br}$ complexes in the wavelength region 390-750 nm are given in fig. 3.4. These cases will now

Figure 3.4: Action spectra for the Na·XCH₃, X=F, Cl, Br, complexes.

The wavelength of the Na D line is given by the dotted line. Vibrational progressions are observed in peaks B and C.

be discussed in turn. Photodepletion is first observed for $\text{Na}\cdots\text{FCH}_3$ at a transition energy corresponding to a shift of about 0.4 eV from the Na D-line (at 589 nm or 2.11 eV) which is the lowest energy transition in the free Na atom[93]. A slightly larger energy shift of about 0.6 eV was found for the lowest electronically-excited state of the $\text{Na}\cdots\text{NH}_3$ complex in a two-colour two-photon ionization experiment by Hertel and co-workers[103]. The shift to lower energies is characteristic of stronger binding in $\text{Na}^*\cdots\text{FCH}_3$ than in $\text{Na}\cdots\text{FCH}_3$.

There are three pronounced photodepletion peaks for $\text{Na}\cdots\text{FCH}_3$ centered at around 485, 590, and 685 nm. According to a previous measurements[19] the peaks were labeled A, B and C respectively. The size of all three peaks were reproduced with an accuracy of $\pm 20\%$ compared with earlier measurements using a different geometry for the molecular beams. The spectral range studied in ref. 19 for photodepletion of $\text{Na}\cdots\text{FCH}_3$ complexes was extended from 450 nm to the present short-wavelength limit of 390 nm. This region of the spectrum showed evidence for an additional weak photodepletion peak, A', centered at 415 nm with a maximum cross-section of $0.02 \text{ \AA}^2$. This section of the spectrum was multiplied by a factor of ten and included in fig. 3.4 as a smoothed spline fit to the experimental data (see insets). The additional fourth peak is clearly the result of excitation to a higher reactive PES. Photodepletion was not observed over the spectral region from 750 to 840 nm for $\text{Na}\cdots\text{FCH}_3$. The assignment of the peaks and their relative intensity will be discussed in terms of *ab initio* theory in the following section.

The action spectrum of $\text{Na}\cdots\text{ClCH}_3$ is shown as the second system in fig.3.4. Although it had not been measured systematically before, a

photodepletion signal was observed at a few wavelength steps between 588 and 593 nm in ref. 17. We observed a broad depletion peak at around 660 nm, labeled C in analogy to the peak for $\text{Na}\cdot\text{FCH}_3$ that appears to the red of the Na D-line. The next peak, B, to shorter wavelengths is centered at around 557 nm. While the peak A is substantial for $\text{Na}\cdot\text{FCH}_3$, though its magnitude is smaller than B and C, peak A for $\text{Na}\cdot\text{ClCH}_3$, centered at approx. 450 nm is just detectable. The ten times magnification of the short wavelength section of the photodepletion spectrum treated with a smoothed spline procedure, indicates a peak cross-section of approx. $0.05 \text{ \AA}^2$. An increase of the depletion cross-section below 410 nm, labeled A', indicates that the next higher PES may also be accessed in this complex.

Photodepletion measurements for $\text{Na}\cdot\text{BrCH}_3$, which is the third system included in fig. 3.4, have not been reported before. The action spectrum shows three depletion peaks. The peak at longest wavelength with a maximum cross-section of $0.95 \text{ \AA}^2$ at around 640 nm, was labeled C. A fairly constant depletion signal of about $0.18 \text{ \AA}^2$ extends to the red, above 700 nm. The next peak toward shorter wavelengths, labeled B, appears at around 565 nm with a maximum cross-section of $1.5 \text{ \AA}^2$. Additionally, a broad peak A was observed at around 480 nm with a maximum cross-section of $0.3 \text{ \AA}^2$.

The action spectra for the peak A in the $\text{Na}\cdot\text{BrCH}_3$ spectrum between 460 and 510 nm as originally recorded, exhibited a progression of superimposed sharp peaks towards negative cross-sections. These peaks were shown to be due to the excitation laser, L_1 , alone. The wavelength and energetic spacing of the spikes corresponded to the

electronic-vibrational excitation, $\text{Na}_2 \rightarrow \text{Na}_2^*(\text{B})$, which was measured by resonant one-color two-photon ionization in ref. 104. The process responsible for enhancing the $\text{Na}\cdots\text{BrCH}_3^+$ complexes at these wavelengths is thought to be $\text{Na}_2^+ + \text{Na}\cdots\text{BrCH}_3 \rightarrow \text{Na}_2 + \text{Na}\cdots\text{BrCH}_3^+$. These peaks were corrected for by substituting the difference between the signal intensities when only the excitation laser (L_1) was fired from when both L_1 and the ionization laser (L_2) were used, for the depletion signal intensity, N , in the Beer-Lambert expression (see section 2.2). We have found no evidence for similar effects involving Na_2^+ , in the wavelength region above the threshold for resonant two-photon ionization of Na_2 at 517 nm (described in ref. 105) or with methyl fluoride or methyl chloride complexes.

The peaks in the action spectrum will be associated with different electronically-excited states of the complex in section 3.4. It is informative to compare the separation between them. Considering the position of both peaks, the location of B and C is more or less symmetric with respect to the Na D-line for $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$. The case of $\text{Na}\cdots\text{FCH}_3$ constitutes a striking anomaly; in this instance the maximum of peak B falls exactly at the Na D-line as if Na were not perturbed by the nearby presence of FCH_3 . Peak B for $\text{Na}\cdots\text{FCH}_3$ is also anomalous in being broader in energy than the corresponding peaks for $\text{Na}\cdots\text{ClCH}_3$ or $\text{Na}\cdots\text{BrCH}_3$. A further noteworthy feature is the progressive increase in breadth of peak C down the series $\text{Na}\cdots\text{XCH}_3$, $\text{X}=\text{F}, \text{Cl}, \text{Br}$. In section 3.4 we shall attribute increasing breadth to the presence of two overlapping excited states contributing to peak C, which become separated in energy down the series $\text{X}=\text{F}, \text{Cl}, \text{Br}$. The intensity of the peaks in the action

spectra will be related to the oscillator strength for the electronic transition and the probability of depletion from the electronically-excited state. Peak A diminishes in going from $X=F$ to Cl and then increases markedly in going from $X=Cl$ to Br .

A reproducible structure is superimposed on the broad peaks B and C for all complexes in fig. 3.4. This structure, which is believed to be vibrational, was not resolved in the measurements with the previous experimental setup[19]. The excitation of stretching and bending modes for the lowest electronically-excited state of $Na\cdots NH_3$ was observed in ref. 103. The linewidth of the vibrational lines in the action spectra for $Na\cdots FCH_3$ indicate a lifetime for the excited-state complexes which cannot be shorter than ~ 0.1 ps. Schulz et al have measured a lifetime of more than 1 ns in a femtosecond pump-probe laser experiment for $Na\cdots NH_3$ [106]. Although, fragmentation of the excited complex was assumed to be the underlying process in their study. The first identified vibrational spacing (v-number unknown) is listed in table 3.1.

In the case of peak C for $Na\cdots FCH_3$, fig. 3.4 shows two progressions with six lines with an energy spacing of 235 cm^{-1} and 241 cm^{-1} . These two progressions are shifted away from one another by an average value of $\sim 90\text{ cm}^{-1}$, indicative of two overlapping progressions. In previous work[17], photodepletion measurements for the complex $Na\cdots FPh$ produced a peak at 635nm (this peak at the red limit of the spectrum, corresponds to C in our terminology) with structure indicating a very similar vibrational spacing of about 240 cm^{-1} . This result indicates that this vibrational mode is likely to arise from stretching of the vdW bond

Table 3.1: The spacing between the first vibrational transitions observed in each progression.

	Na··FCH ₃	Na··ClCH ₃	Na··BrCH ₃
c	235cm ⁻¹	207cm ⁻¹	182cm ⁻¹
c'	241cm ⁻¹		129cm ⁻¹
c''			124cm ⁻¹
b	91cm ⁻¹	142cm ⁻¹	145cm ⁻¹

in either the ground or excited state of the complex which should not be strongly dependent on the radical, $R = \text{CH}_3$ or Ph.

Within a given progression, a Birge-Sponer plot for ΔE_v (illustrated in fig. 3.5) is approximately linear. For peak C of $\text{Na}\cdots\text{FCH}_3$ the anharmonicity constants, taken from the slopes of the plots in fig. 3.5, are 11 ± 2 and $13 \pm 1 \text{ cm}^{-1}$ respectively. The anharmonicity constant for each observed progression is listed in table 3.2.

A trend of decreasing vibrational level spacing and in general decreasing anharmonicity constant was observed for peak C from $X=\text{F}$ to Cl to Br . In case of $\text{Na}\cdots\text{ClCH}_3$, the anharmonicity constant was only calculated from the first to the eighth vibrational spacing. There appeared to be an irregularity in the vibrational progression after the ninth line (see fig. 3.5). Peak C of $\text{Na}\cdots\text{BrCH}_3$ carries two vibrational progressions. We are able to identify at least 9 lines for each progression. These two progressions are shifted from one another by an average value of $64 \pm 6.5 \text{ cm}^{-1}$.

For peak B the vibrational level spacing and anharmonicity constant for the systems $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ are very similar. However, peak B for $\text{Na}\cdots\text{FCH}_3$ was observed to have anomalous properties in regard to both vibrational level spacing and anharmonicity constant as compared to the other two systems.

We are not able to identify vibrational structure for the other peaks A or A', with the exception of peak A in the $\text{Na}\cdots\text{FCH}_3$ action spectrum. The vibrational structure for peak A in fig. 3.4 for $\text{Na}\cdots\text{FCH}_3$, which was not present under previous experimental conditions, was

Figure 3.5: Birge-Sponer plots for the vibrational progressions observed in the experimental action spectra in fig. 3.4. The appropriate vibrational quantum number for each line (v') is given relative to that for the first line (v_0) observed in each progression.

Table 3.2: The anharmonicity constant observed in each vibration progression.

	Na··FCH ₃	Na··ClCH ₃	Na··BrCH ₃
c	$11 \pm 2\text{cm}^{-1}$	$7.0 \pm 0.5\text{cm}^{-1}$	$13 \pm 3\text{cm}^{-1}$
c'	$13 \pm 1\text{cm}^{-1}$		$5 \pm 2\text{cm}^{-1}$
c''			$4.0 \pm 0.5\text{cm}^{-1}$
b	$1.5 \pm 0.5\text{cm}^{-1}$	$22 \pm 5\text{cm}^{-1}$	$18 \pm 1\text{cm}^{-1}$

identified with the vibrational structure of the $\text{Na}_2(\text{B})$ state. Since the TOF spectrum clearly distinguishes between the Na_2^+ and $\text{Na}\cdots\text{FCH}_3^+$ ion signals, the structure must arise from reaction between neutral $\text{Na}\cdots\text{FCH}_3$ complexes and sodium-dimer molecules. We attribute the depletion to collisions between Na_2^* and $\text{Na}\cdots\text{FCH}_3$. The Na_2^* is only present at wavelengths corresponding to resonant two-photon ionization of Na_2 [104]. The depletion process would be collisionally induced reaction; $\text{Na}_2^* + \text{Na}\cdots\text{FCH}_3 \rightarrow \text{Na}_2 + \text{NaF} + \text{CH}_3$, $\Delta H^\circ = -11 \text{ kcal/mol}$ (using $D^0(\text{F-CH}_3)=113 \text{ kcal/mol}$ [93] and $D^0(\text{NaF})=124 \text{ kcal/mol}$ [93]).

3.4 Discussion

peak assignment

An assignment of the electronically-excited states of the complex, $\text{Na}^*\cdots\text{XCH}_3$, will be based on the atomic states of Na which are perturbed by the XCH_3 molecule. Electron-density contour plots, obtained by *ab initio* calculation[96], for the highest occupied molecular orbital (HOMO) in the ground and electronically-excited states of the $\text{Na}\cdots\text{FCH}_3$ complex are illustrated in fig 3.6. The character of the electron distribution around the Na atom clearly resembles distinct atomic orbitals. Each contour plot has been labeled with the appropriate atomic-term symbol which arises from the orbital occupation on the Na atom. The orientation of the Na 3p orbitals can give rise to either a σ interaction from the p_z orbital, or a π interaction from the p_x and p_y orbitals, with the closed-shell XCH_3 molecule. The subscripts x, y and z have been appended to the atomic term symbol to indicate the alignment of the p orbital. The molecular orbital formed from the Na p_y orbital cannot be

Figure 3.6: Calculated electron-density distribution for the HOMO in the ground ($3s^2S$) and the excited ($3p^2P_x$, $3p^2P_z$ and $4s^2S$) states of $\text{Na}\cdot\text{FCH}_3$ [96]. The broken line indicates where the electron density is zero, while the solid and dotted lines indicate positive and negative-phase values for the electron density. The increment is $0.0025/\text{\AA}^2$. The plots were made by MOLDEN.

visualized in the plane of the complex shown in fig.3.6, but it resembles the HOMO for the $3p\ 2P_x$ excited state closely. The *ab initio* calculation, performed by others in this group, is described in detail in ref. 96. The features of this calculation, which have enabled a better understanding of the experimental spectra, will be described in detail in the following paragraphs.

The excited states of the three complexes were studied at the configuration interaction (CI) level incorporating single excitations only. The CI-single (CIS) method was encoded into Gaussian 92. The basis set used for each atom was described in section 3.2. The neglect of double (and higher) excitations lead to a significant quantitative uncertainty in these calculations but they should be sufficient for a qualitative description of the excited states and an assignment of the peaks observed in the experimental depletion spectrum.

The potential energy of the electronically-excited states of the complex were first calculated with the nuclei held fixed in the ground-state geometry described in fig. 3.3. The calculated potential energy will therefore be equivalent to a vertical-excitation energy. It is expected that the complexes formed in the crossed-molecular beams will have considerable population of vibrationally-excited levels. Therefore, the onset for absorption is likely to be observed from high vibrational levels of the ground-state complex to low vibrational levels in the excited-state complex. Therefore, as a rough guide, the potential energy of the electronically-excited states relative to the ground-state dissociation continuum will be compared to the onset for photodepletion in the experimental spectrum. The comparison is made in table 3.3 for the

Table 3.3: Location of peaks in the action spectrum and the calculated vertical transition energies[96] to lowest excited states for $\text{Na}\cdots\text{FCH}_3$.

Photodepletion peak	C	B	A	A'
<i>Excitation energy¹ / eV</i>	1.69	1.97	2.41	2.82
Electronic state	$^2\text{P}_x / ^2\text{P}_y$	$^2\text{P}_z$	^2S	^2D
<i>MRDCI² / eV</i>	1.50/1.56	2.10	2.37	2.83
<i>CIS² / eV</i>	1.43 / 1.46	2.07	2.29	2.70

¹corresponding to the onset for photodepletion in the experimental spectrum.

²the potential energy of the electronic-excited complex has been calculated relative to the ground state energy of $\text{Na}+\text{FCH}_3$ using MRDCI and CIS methods.

Na $\cdot$ FCH₃ complex. For this system, the CIS calculation of the vertical transition energy is supplemented by a more extensive treatment of electron correlation using a multi-reference calculation with single and double excitations (MRDCI)[96].

The four broad peaks which appear in the experimental photodepletion spectra of fig. 3.4 were assigned according to the calculated sequence of excited states. The broad peak C is assigned to the superposition of the depletion cross-section in which the complexes were excited to the 3p 2P_x and 2P_y states, peak B corresponds to excitation to the 3p 2P_z state, peak A is a measure of the photodepletion by way of the normally forbidden transition to the 4s 2S state, and peak A' is photodepletion by excitation to the 3d 2D state. The following paragraphs enlarge on these assignments.

The electronically-excited states of the complex defined above are the TS, i.e. [Na $\cdot$ XCH₃] ‡ , for the reaction Na $\cdot$ +XCH₃ $\rightarrow$ NaX+CH₃. Therefore, the assignment of the peaks measured in the action spectra in fig.3.4, can be interpreted as representing the appropriate atomic state of Na $\cdot$ which correlates with the TS formed by photoexcitation of the ground state complex, Na $\cdot$ XCH₃.

The position of each broad peak in the experimental action spectra corresponds to the location of electronically-excited states of the complex. However, the breadth of each peak does not necessarily correspond to the range of possible wavelengths for which the Franck-Condon factors for excitation are finite, but gives the range for which reaction occurs, resulting in depletion of the complex. If excitation

occurs to excited-state configurations from which a reaction pathway is not accessible, then it will not correspond to an absorption feature in the experimental spectrum. This might explain the discrepancy with the onset for photodepletion for the $3p\ ^2P_{x,y}$ state of $\text{Na}^+\cdots\text{FCH}_3$ in the experimental action spectrum and the calculated vertical-transition energy (see table 3.3). Excitation to low vibrational levels of the $3p\ ^2P_{x,y}$ state of $\text{Na}^+\cdots\text{FCH}_3$ might not lead to reaction of the complex and the depletion cross section in the action spectrum will then be zero. Depletion is only observed when excitation occurs to higher vibrational levels. This observation will be shown in chapter 5, to be in accord with experiments in which the lifetime of the excited state was measured. In the latter work, excited-state complexes, $\text{Na}^+\cdots\text{FCH}_3$, were detected using low excitation energies at which photodepletion was not observed.

There is a significant stabilization of the $4s\ ^2S$ and $3d\ ^2D$ states in the complex relative to the atomic states of Na. The $3s\ ^2S \rightarrow 4s\ ^2S$ and $3s\ ^2S \rightarrow 3d\ ^2D$ transitions in atomic Na occur at energies of 3.20 and 3.63eV above the $\text{Na} + \text{FCH}_3$ dissociation continuum[93]. The experimental photodepletion spectra and the MRDCI calculations both estimate that the $4s\ ^2S$ and $3d\ ^2D$ states of the $\text{Na}\cdots\text{FCH}_3$ complex exist at energies of ~2.4 and ~2.8eV.

An assignment of the peaks in the action spectra for $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ is shown in tables 3.4 & 3.5 based on calculations of the potential energy for the excited-state complexes calculated at the CIS level using Gaussian92. If the calculated vertical-transition energy is greater than the excitation energy for the corresponding transition in atomic Na, then it must represent excitation to a state which is

Table 3.4: Location of peaks in the action spectrum and the calculated vertical transition energies[96] to lowest excited states for $\text{Na}\cdot\cdot\text{ClCH}_3$.

Photodepletion peak	C	B	A	A'
<i>Excitation energy¹ / eV</i>	1.66	2.13	2.63	3.00
Electronic state	$^2\text{P}_x / ^2\text{P}_y$	$^2\text{P}_z$	^2S	^2D
<i>Vertical transition energy² / eV</i>	1.7/1.8	2.3	2.8	3.2

¹corresponding to the onset for photodepletion in the experimental spectrum.

²the potential energy of the electronic-excited complex has been calculated relative to the ground state energy of $\text{Na}+\text{ClCH}_3$ using Gaussian92 at the CIS level.

Table 3.5: Location of peaks in the action spectrum and the calculated vertical transition energies[96] to lowest excited states for $\text{Na}\cdots\text{BrCH}_3$.

Photodepletion peak	C	B	A	A'
<i>Excitation energy¹ / eV</i>	1.86	2.11	2.34	
Electronic state	$^2\text{P}_x / ^2\text{P}_y$	$^2\text{P}_z$	^2S	^2D
<i>Vertical transition energy² / eV</i>	1.8/1.9	2.3	2.9	3.3

¹corresponding to the onset for photodepletion in the experimental spectrum.

²the potential energy of the electronic-excited complex has been calculated relative to the ground state energy of $\text{Na}+\text{BrCH}_3$ using Gaussian92 at the CIS level.

dissociative along the Na-X coordinate. The $3p\ ^2P$ excited state for atomic Na is located at an excitation energy of 2.11 eV. Therefore, for both $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$, a vertical transition accessed a bound region of the $3p\ ^2P_{x,y}$ state, while a repulsive region of the $3p\ ^2P_z$ state was reached. When the complexes are excited into repulsive states with respect to the Na-X coordinate, their van der Waals bonds can be expected to break.

Qualitative profiles for the potential energy of the three complexes which are consistent with the experimental photodepletion spectra reported here and the *ab initio* calculations[96] are shown in fig. 3.7. The van der Waals interaction between Na and XCH_3 induces a splitting between the states $3p\ ^2P_z$ and $3p\ ^2P_{x,y}$. The interaction between the paired electrons in the orbitals of the closed-shell halide, XCH_3 , and the empty p_σ orbital of $\text{Na}^*(3p\ ^2P_{x,y})$ gives rise to stabilization of the complex. However, the interaction becomes repulsive in $\text{Na}^*(3p\ ^2P_z)$ due to unfavourable overlap between the occupied $3p_z$ orbital and the closed-shell halide molecule. This repulsion is not sufficiently compensated for by attractive dispersion interactions in $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$, and the $3p\ ^2P_z$ state remains unbound. The energy splitting is clearly observed in the experimental photodepletion spectra of fig.3.4 for the systems $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$, in which peaks B and C are located to either side of the D-line. For the case of $\text{Na}\cdots\text{FCH}_3$, the experimentally measured peak B was found to be located at the sodium D-line. The anomalous position of peak B for this system can be explained by the different nature of the $3p\ ^2P_z$ state for $\text{Na}\cdots\text{FCH}_3$ which was determined by *ab initio* calculation to be a bound state (see ref. 96). In this case,

Figure 3.7: Qualitative potential-energy profiles referring to the $\text{Na} \cdots \text{X}$ bond distance for the three complexes. The well depths for the ground-state complexes were taken from *ab initio* calculations[96] but the excited states were adjusted to match experimental transition energies represented by arrows. The $3p \ ^2P_x$ state is similar to the $3p \ ^2P_y$ state which is not shown.

attractive dispersion interactions in $\text{Na}^+\cdots\text{FCH}_3$ are sufficient to stabilize the excited-state complex.

Although the potential-energy curves in fig. 3.7 are consistent with the location of the electronically-excited states of the complex and the peaks observed in the photodepletion spectra, they do not indicate the dynamics of the reaction which follows excitation of $\text{Na}\cdots\text{XCH}_3$. The excited-state complex is the TS for the reaction $\text{Na}^+\text{XCH}_3 \rightarrow \text{NaX}+\text{CH}_3$ and the dissociation of the complex to give products is not described in fig. 3.7. However, the evolution of the TS will be described for a similar system, $\text{Na}^+\text{FH} \rightarrow \text{NaF}+\text{H}$ in chapter 4 based on a more detailed *ab initio* calculation of the complete PES for the ground and excited states of NaFH .

A further splitting between the $3p\ ^2P_x$ and $3p\ ^2P_y$ states will be induced in bent configurations of the complexes by the CH_3 umbrella head. This effect was not resolved in the action spectra in fig. 3.4 which involve excitation of $\text{Na}\cdots\text{XCH}_3$ from (minimum-energy) bent configurations of the bound complex. However, the splitting might be the origin of the considerable breadth of peak C relative to peak B in fig3.4(b) for $\text{Na}\cdots\text{ClCH}_3$ and the presence of two vibrational progressions within peak C in fig. 3.4(a),(c) for $\text{Na}\cdots\text{FCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$. In the latter case, each progression belonging to a different electronically-excited state.

peak intensity

The depletion cross-section is a product of the cross section for absorption of a photon of wavelength λ_1 by the ground-state complex, and the probability of removal following excitation which leads to

photodepletion. The oscillator strength for each electronic transition, f_i , has been determined at the CIS level using the program 'Gaussian92'[96]. The cross-section for photoabsorption integrated across the appropriate energy interval for each electronically-excited state, i , is calculated from

$$\int \sigma_{\text{abs}} dE = \frac{\alpha h^2}{2m_e} \sum_i f_i \quad 3.1$$

where α is the unitless fine-structure constant ($\alpha=1/137$), h is Boltzmann's constant and m_e is the mass of an electron[109]. The depletion cross-section has been integrated across the energy interval for each peak in the action spectra in fig.3.4. The comparison between the (measured) depletion and the (calculated) absorption cross-section is made for the $\text{Na}\cdots\text{FCH}_3$ complex in table 3.6. The depletion probability ($\rightarrow\text{NaF}+\text{CH}_3$ of $\text{Na}+\text{FCH}_3$) following excitation to each electronic state is the ratio of the two values for the cross-section. There is an increasing probability of depletion with increasing electronic excitation of the complex; i.e. from ~10% for the $3p\ ^2P$ state to ~100% for the $3d\ ^2D$ state. This result is in accord with an increasing reaction cross-section observed experimentally by Lee and co-workers for the direct reaction between Na^* and BrCH_3 measured in crossed-molecular beams[49].

According to the transition-selection rule ($\Delta L=\pm 1$), for a free Na atom the transitions from the ground state, $3s\ ^2S$, to the higher energy states, $4s\ ^2S$ and $3d\ ^2D$, are forbidden (the oscillator strength for these electronic transitions in atomic Na are $\sim 10^{-5}$ [93]). The HOMO for the $4s\ ^2S$ state of Na perturbed by FCH_3 was calculated to consist mainly of the $4s$ orbital on Na with a small portion of the p_z orbital mixed in, as is indicated in fig.3.6. As a result, in the environment of the complex, these

Table 3.6: Integrated depletion cross-section from the Na·FCH₃ action spectrum and the calculated oscillator strength[96] for each electronic transition.

Photodepletion peak	C	B	A	A'
<i>Integrated depletion cross-section /Å²eV</i>	0.061	0.105	0.026	0.003
Electronic state	² P _x + ² P _y	² P _z	² S	² D
<i>Oscillator strength</i>	0.3+0.3	0.3	0.06	0.003
<i>Integrated absorption cross-section /Å²eV</i>	0.7	0.3	0.07	0.003
<i>Depletion probability</i>	10%	30%	40%	~100%

transitions become more allowed. However, the relative intensity of peak A in the photodepletion spectra also appears to arise from an increased probability of depletion from the $4s\ ^2S$ state (see table 3.6).

The probability of depletion from each electronically-excited state has been determined for $\text{Na}\cdot\cdot\text{ClCH}_3$ and $\text{Na}\cdot\cdot\text{BrCH}_3$ in tables 3.7 & 3.8. The depletion probability was again observed to increase with increasing electronic excitation, i.e. $3p\ ^2P_{x,y} < 3p\ ^2P_z < 4s\ ^2S$. In addition, for a particular excited state the depletion probability increases for the series of complexes $\text{Na}\cdot\cdot\text{XCH}_3$ in the order $\text{X} = \text{F} < \text{Cl} < \text{Br}$. Molecular-beam experiments have shown an increase of $\sim 2\times$ in the reaction cross-section in collisions between $\text{Na}^*(4s) + \text{BrCH}_3$ as compared with $\text{Na}^*(3p) + \text{BrCH}_3$ [49]. However, the experimentally observed substantial depletion cross-section for the $4s\ ^2S$ excited state of $\text{Na}\cdot\cdot\text{BrCH}_3$ cannot be solely explained by a higher depletion probability for $\text{Na}\cdot\cdot\text{BrCH}_3$ in going from $3p\ ^2P$ (peak B and C) to $4s\ ^2S$ (peak A) as the excited state since the integrated depletion cross-section of $\text{Na}\cdot\cdot\text{BrCH}_3$ is much larger than calculated[96] absorption cross section.

A further contributing factor, which could increase the $3s\ ^2S \rightarrow 4s\ ^2S$ oscillator strength, is the effect of the presence of the large Br nucleus on the spin and angular momentum selection rules. The effect of spin-orbit coupling was not incorporated into the CIS calculation. For $\text{Na}\cdot\cdot\text{BrCH}_3$, only the total electron angular momentum selection rule ($\Delta J=0, 1$) is a good one, whereas in both the $3s\ ^2S$ and $4s\ ^2S$ states, atomic Na has a total electronic angular momentum, J , equal to $1/2$.

Table 3.7: Integrated depletion cross-section from the Na $\cdot$ -ClCH₃ action spectrum and the calculated oscillator strength[96] for each electronic transition.

Photodepletion peak	C	B	A	A'
<i>Integrated depletion cross-section / Å²eV</i>	0.171	0.103	0.013	
Electronic state	² P _x + ² P _y	² P _z	² S	² D
<i>Oscillator strength</i>	0.3+0.3	0.4	0.01	0.001
<i>Integrated absorption cross-section / Å²eV</i>	0.7	0.4	0.01	0.001
<i>Depletion probability</i>	20%	30%	~100%	

Table 3.8: Integrated photodepletion cross-section from the Na $\cdots$ BrCH₃ action spectrum and the calculated oscillator strength[96] for each electronic transition.

Photodepletion peak	C	B	A	A'
<i>Integrated depletion cross-section / Å²eV</i>	0.17	0.17	0.12	
Electronic state	² P _x + ² P _y	² P _z	² S	² D
<i>Oscillator strength</i>	0.3+0.3	0.4	0.01	0.001
<i>Integrated absorption cross-section / Å²eV</i>	0.7	0.4	0.01	0.001
<i>Depletion probability</i>	20%	40%		

vibrational structure

In this section we discuss the structure which was found for the three complexes; $\text{Na}\cdot\text{XCH}_3$, $\text{X} = \text{F}, \text{Cl}$ and Br , in the experimental action spectra in fig 3.4 with respect to vibrational frequencies determined from *ab initio* calculation[96].

Figure 3.8 compares the minimum-energy structures for the three lowest excited states, $3p\ ^2P_x$, 2P_y and 2P_z , of the complex $\text{Na}\cdot\text{FCH}_3$ obtained at the CIS level with the ground-state structure of the complex calculated at the Moller-Plesset level (see section 3.2 and ref. 96). Compared to the geometry of the ground state, the three excited states have a reduced $\text{Na}\cdot\text{F}$ internuclear distance and nearly linear structure. Thus, for a vertical excitation, it is expected that $\text{Na}\cdot\text{F}$ stretching modes and $\text{Na}\cdot\text{F-C}$ bending modes will both be excited. In addition, stretching and bending modes are likely to be populated initially in the ground-state complex which is formed in the crossed-molecular beams. Therefore, the progressions observed in the experimental action spectra should be compared with both ground and excited-state vibrational frequencies.

The vibrational frequencies calculated for the $\text{Na}\cdot\text{FCH}_3$ complex in the ground and excited states are listed in table 3.9. In section 3.3 the vibrational structure observed in peak C for $\text{Na}\cdot\text{FCH}_3$ (see fig 3.4) was identified as two progressions. The two progressions were shifted from each other by an average energy of $\sim 90\text{cm}^{-1}$, which is the same order of magnitude as the calculated energy splitting for the two states, namely $\sim 0.01\text{eV}$ (see table 3.3). The two progressions may, therefore, be carried by

Excited State: $3p\ ^2P_z$ Dissociation
energy

0.13 eV

 $3p\ ^2P_{x,y}$

0.51 eV

Ground State: $3s\ ^2S$

0.21 eV

Figure 3.8: Geometry and dissociation energy of the most stable configurations for the excited states of Na $\cdot$ FCH₃ calculated using Gaussian92 at the CIS level[96]. The structure of the ground-state complex has been reproduced from fig. 3.3 for comparison.

Table 3.9: Calculated frequencies for vibrational modes in ground¹ and excited² states of Na··FCH₃[96].

Electronic state	3s ² S	3p ² P _x	3p ² P _y	3p ² P _z
Na-F stretch ³ /cm ⁻¹	178	228	227	282
Na-F-C bend /cm ⁻¹	78	63	47	110

¹using Gaussian92 at the MP2 level.

²using Gaussian92 at the CIS level.

³the stretching frequency is 536cm⁻¹ in the NaF molecule[98].

the two excited states which are identified as $3p\ ^2P_x$ and $3p\ ^2P_y$. However, another possible explanation for the origin of the second progression is the excitation of one quantum of the in-plane inversion mode of $\text{Na}\cdots\text{FCH}_3$ with a calculated frequency of 63 and 47cm^{-1} for $3p\ ^2P_x$ and $3p\ ^2P_y$ respectively. The spacing between vibrational transitions within each progression corresponds to the calculated vibrational frequencies for the $\text{Na}\cdots\text{F}$ stretching mode in these two excited states. The values obtained from the experimental spectrum in fig. 3.4 are 235 cm^{-1} and 241 cm^{-1} , and the calculated frequencies are 228 and 227 cm^{-1} for the $3p\ ^2P_x$ state and the $3p\ ^2P_y$ state respectively.

A single vibrational progression was observed in peak B for $\text{Na}\cdots\text{FCH}_3$ with a much shorter spacing of approx. 91 cm^{-1} . The $3p\ ^2P_z$ state for $\text{Na}\cdots\text{FCH}_3$ was calculated to be a bound state with a frequency for the $\text{Na}\cdots\text{F}$ stretching mode of 282 cm^{-1} and for the bending mode of 110 cm^{-1} . In comparison with the experimental value 91cm^{-1} we assume for the case of the $3p\ ^2P_z$ state that the bending mode was excited.

For the system $\text{Na}\cdots\text{FCH}_3$, the minimum-energy geometry of the 3^2P_z state is very different from that for the $3p\ ^2P_{x,y}$ states, with a shorter $\text{F}\cdots\text{Na}$ distance and a Na-F-C angle close to 180° . In the case of $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ complexes, only the $3p\ ^2P_{x,y}$ states were calculated to be bound while the $3p\ ^2P_z$ state was found to be dissociative along the Na-X coordinate. The geometry of the excited states for these two systems are illustrated in fig. 3.9. The frequencies of the vibrational modes in the ground and excited states are given in tables 3.10 & 3.11. The frequencies for the Na-X stretching mode and the Na-X-C bending mode decrease in the order $\text{X}=\text{F} > \text{Cl} > \text{Br}$.

Excited State: $3p\ ^2P_z$
unbound

Excited State: $3p\ ^2P_x\ (^2P_y)$

Ground State: $3s\ ^2S$

Figure 3.9: Geometry and dissociation energy (D) of the most stable configurations for $\text{Na}\cdots\text{XCH}_3$, $\text{X}=\text{Cl}, \text{Br}$, calculated using Gaussian92 at the CIS level[96]. The structure of the ground-state complexes have been reproduced from fig. 3.3 for comparison.

Table 3.10: Calculated frequencies for vibrational modes in ground¹ and excited² states of Na··ClCH₃[96].

Electronic state	3s ² S	3p ² P _x	3p ² P _y
Na-Cl stretch ³ /cm ⁻¹	79	156	126
Na-Cl-C bend /cm ⁻¹	55	113	78

¹using Gaussian92 at the MP2 level.

²using Gaussian92 at the CIS level.

³the stretching frequency is 365cm⁻¹ in the NaCl molecule[98].

Table 3.11: Calculated frequencies for vibrational modes in ground¹ and excited² states of Na··BrCH₃[96].

Electronic state	3s ² S	3p ² P _x	3p ² P _y
<i>Na-Br stretch</i> ³ <i>/cm⁻¹</i>	17	151	109
<i>Na-Br-C bend</i> <i>/cm⁻¹</i>	1	96	62

¹using Gaussian92 at the MP2 level.

²using Gaussian92 at the CIS level.

³the stretching frequency is 298cm⁻¹ in the NaBr molecule[98].

A single progression in peak C was identified for $\text{Na}\cdots\text{ClCH}_3$ in the experimental action spectrum (see fig. 3.4). The calculated splitting between the $3p\ ^2P_x$ and 2P_y states for $\text{Na}\cdots\text{ClCH}_3$ is much larger than the expected vibrational frequency for the Na-Cl stretching mode. The considerable width of the vibrational lines could include a superposition of two progressions. Therefore the first lines to the red side of the progression are due to the lowest excited state, $3p\ ^2P_x$, whereas the last lines to the blue side are very likely of $3p\ ^2P_y$ origin. An irregularity in the vibrational spacing after the ninth line of the progression supports this model (see the Birge-Sponer plots in fig.3.5). The calculated excited-state vibrational frequency for the $\text{Na}\cdots\text{Cl}$ stretching mode of the lowest excited state was 156 cm^{-1} , which corresponds approximately with the spacing between the first two vibrational lines of the progression, namely 207 cm^{-1} . The calculated vibrational frequency in the second excited state, $3p\ ^2P_y$, was 126 cm^{-1} .

The vibrational structure of peak C for $\text{Na}\cdots\text{BrCH}_3$ consisted of two vibrational progressions, with a spacing of approx. 129 and 124 cm^{-1} . Again the two progressions may belong to the two excited states, $3p\ ^2P_x$ and 2P_y , for which the $\text{Na}\cdots\text{Br}$ stretching frequencies were 151 cm^{-1} and 109 cm^{-1} respectively. However, it is also possible that the second progression originates in the excitation of one quantum of the in-plane inversion mode of $\text{Na}\cdots\text{BrCH}_3$ with a calculated frequency of 62 cm^{-1} .

A single vibrational progression was observed in peak B for $\text{Na}\cdots\text{XCH}_3$, $\text{X}=\text{Cl}$ and Br with a spacing of approx. 142 cm^{-1} , and 145 cm^{-1} respectively. These vibrational spacings are much shorter compared to those in peak C. The $3p\ ^2P_z$ state for $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ were

identified to be repulsive. Excitation to a repulsive state would be expected to yield a broad unstructured peak. The presence of vibrational structure in peak B for these two systems could, however, be due to binding along other degrees of freedom than that shown in the qualitative potential energy curves in fig. 3.7. Similar observation of vibrational modes perpendicular to the dissociation coordinate in a repulsive state has been made by Newmark and co-workers[28,29] in their Transition State Spectroscopy studies using negative-ion photodetachment.

3.5 Summary

The van der Waals (vdW) complexes $\text{Na}\cdots\text{XCH}_3$, $\text{X} = \text{F}, \text{Cl}$ and Br , were observed in the crossed-beam apparatus. In other work from this laboratory, the complex $\text{Na}\cdots\text{FCH}_3$ was determined by means of *ab initio* calculation[96] to be the most stable of the series. A bond energy of 0.21 eV was calculated in comparison to ~ 0.04 eV for $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$. This result was in accord with experiment in which a much higher concentration of these complexes was obtained.

Reaction of the complex, in the experiments presented in this thesis, was found to be initiated by photoexcitation across a broad range of wavelengths. The depletion cross-section was measured in the wavelength region from 390 to 750 nm. Pronounced maxima were found at short wavelengths and to either side of the Na D-line. The initial excitation occurred to a selected Transition State (TS) configuration on a potential-energy surface (PES) which would be traversed by the reaction $\text{Na}^* + \text{XCH}_3 \rightarrow \text{NaX} + \text{CH}_3$, in which the reactant alkali atom is in an

electronically-excited state. Photoexcitation within the above wavelength range was found to initiate the reaction between XCH_3 and electronically-excited Na in the $3p\ ^2P_{x,y}$, $3p\ ^2P_z$, $4s\ ^2S$ and $3d\ ^2D$ excited states. The photodepletion peaks were assigned according to the atomic state of Na^* which correlates with the TS formed by excitation of $\text{Na}\cdot\text{XCH}_3$. This assignment was assisted by *ab initio* calculations of the transition energies for electronic excitation of the complexes[96].

The electronically-excited states of the complexes determined by *ab initio* calculation were mainly characterized by the atomic state of the Na atom in the asymptotic limit. However, excited states were observed in the action spectra for the complexes which corresponded to the forbidden transition of an electron from the $3s$ to $4s$ and $3d$ orbitals in the Na atom. This was explained by the admixture of the $3p$ orbital with the $4s$ and $3d$ orbitals when the Na atom is bound in the complex. The atomic transition energies to $4s$ and $3d$ were observed in the experimental action spectrum to be lowered by about 0.6eV for the case of $\text{Na}\cdot\text{FCH}_3$. The van der Waals interaction caused an energy splitting between the $3p\ ^2P_z$ state and the $3p\ ^2P_{x,y}$ state. A much weaker energetic splitting between the $3p\ ^2P_x$ and the $3p\ ^2P_y$ orbital was expected to be induced by the CH_3 umbrella in the bent $\text{Na}\cdot\text{XCH}_3$ complex. Although we found indirect experimental evidence for the predicted splitting, the two states could not be resolved in the photodepletion spectra.

As in comparable Transition State Spectroscopy (TSS) experiments performed by Soep and co-workers on the system $\text{Ca}\cdot\text{XH}$ ($\text{X} = \text{Cl}, \text{Br}$), the TS accessed by photoexcitation of the complex lasts for a sufficient length of time (at least 1ps in ref. 10) to yield vibrational structure. Clear

vibrational progressions were identified in the action spectra within the $3p\ ^2P_{x,y}$ and $3p\ ^2P_z$ photodepletion peaks. In the case of the superimposed $3p\ ^2P_{x,y}$ states the vibrational spacing agreed very well with the calculated frequencies for the stretching mode for the van der Waals bond in the TS[96]. The $3p\ ^2P_z$ potential curve was calculated to be bound along the $\text{Na}\cdots\text{X}$ axis for $\text{X}=\text{F}$, but repulsive for $\text{X}=\text{Cl}$, Br . The measured vibrational structures for $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$ are likely to be caused by oscillations perpendicular to the bond direction. For the case of $\text{Na}\cdots\text{F}$, the vibrational spacing matched the calculated value for the bending mode in the TS, $[\text{Na}\cdots\text{FCH}_3]^\ddagger$.

In chapter 5, the photodepletion spectrum for $\text{Na}\cdots\text{FCH}_3$ will be compared with a different action spectrum for the same system in which the TS is probed directly in the frequency-domain. An estimate of the concentration of TS species in the duration of the excitation-laser pulse has been made. This was achieved via direct ionization of TS species using a wavelength for L_2 which could not ionize $\text{Na}\cdots\text{FCH}_3$. The TS ionization measurements have provided an estimate for the (widely varying) lifetime of the transition state configurations in the harpooning reaction.

In the following chapter (ch.4), a more detailed description of the dynamics will be given for the excited-state reaction Na^*+HF based on an accurate *ab initio* calculation of the complete adiabatic PESs for the ground and electronically-excited states of the system NaFH performed by others in this laboratory[110]. The charge-transfer dissociation reaction, which is likely to closely reflect that observed in $\text{Na}\cdots\text{XCH}_3$ complexes, will be attributed to surface hopping from the excited state to

the ground state into the product channel $\text{NaF}+\text{H}$. This surface-hop event occurs from a largely-covalent configuration in the excited state to a largely-ionic configuration in the ground state, and is, therefore directly related to 'harpooning'.

Chapter 4

Photodepletion of Na·FH

The van der Waals complex Na·FH has been observed for the first time in a crossed-molecular beam apparatus. Experimental results on the photoinduced charge-transfer dissociation reaction of Na·FH will be related to an *ab initio* study of the same process performed by others in this laboratory[110]. The increased simplicity obtained by substituting a H atom for a CH₃ group has enabled more detailed calculations to be performed on the ground and electronically-excited states of this complex which assist in a description of the dynamics which follow excitation of Na·FH.

Photodepletion of the complex, Na·FH, is believed to occur through excitation of the sodium chromophore followed by charge-transfer to the hydrogen-halide molecule, i.e. $\text{Na}\cdot\text{FH} + h\nu_1 \rightarrow [\text{Na}^*\cdot\text{FH}]^\ddagger \rightarrow [\text{Na}^+\cdot\text{FH}^-]^\ddagger \rightarrow \text{NaF} + \text{H}$ (or $\text{Na} + \text{FH}$). Photoexcitation was probed in a broad spectral region, 440-840nm, to either side of the D-line transition (589nm) in atomic Na. The observed action spectrum for photodepletion consisted of two broad peaks with vibrational structure; a weaker peak centered at 780nm and a stronger one at 650nm. *Ab initio* calculations[110] indicated the presence of excited states of the complex, Na*·FH, in the wavelength region corresponding to the observed peaks. These electronically-excited states correspond to Na in the 3p ²P state perturbed by the HF molecule. Extensive *ab initio* calculation[110] of the potential-energy surface (PES) for the ground and excited states of NaFH revealed an avoided-crossing between the first excited state and the ground state which could lead to dissociation of the excited-state

complex ($\rightarrow \text{NaF}+\text{H}$) from a restricted range of transition states configurations.

4.1 Identification of $\text{Na}\cdots\text{FH}$ complexes

The van der Waals complex $\text{Na}\cdots\text{FH}$ was generated by crossing the Na atomic beam with an expansion of a mixture of HF in He from the pulsed-valve source at a backing pressure of $\sim 200\text{kPa}$. The concentration of complexes was measured by photoionization at $\lambda_2=248\text{nm}$. The complex-ion signal was measured using different concentrations of HF in both He and Ar carrier gas. It was found that the intensity of complexes decreased if the fraction of HF rose above 2.5% and also when Ar was used as the carrier gas. This effect is illustrated in fig.4.1. These conditions were in contrast to those which were optimum for generating $\text{Na}\cdots\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) complexes. For $\text{Na}\cdots\text{FCH}_3$ a higher concentration of the halide was used (although the exact composition of the expansion mixture was uncertain- see section 2.1), with Ar as the carrier gas, at a much higher backing pressure for the pulsed-valve source (400kPa).

A likely reason for requiring a lower concentration of HF in the expansion mixture is due to increased association of the highly polar HF molecules to give clusters, $(\text{HF})_n$, with $n > 2$, as the concentration of HF is increased. In addition, when Ar was used as the carrier gas even larger clusters, $(\text{Ar})_n\cdots(\text{HF})_n$, will be present in the expansion. However, a He atom will bind with a HF molecule more weakly due to its lower polarizability, and the lower backing pressure used will further reduce the size of $(\text{HF})_n$ clusters formed in the expansion. The presence of large clusters appears to inhibit the formation of the complex, $\text{Na}\cdots\text{FH}$, in the

Figure 4.1: The dependence of the complex-ion signal for $\text{Na}\cdots\text{FH}$ on the composition and identity of the carrier gas in the expansion mixture.

crossed-molecular beams.

The most probable source of $\text{Na}\cdots\text{FH}$ complexes is the exchange reaction $\text{Na} + (\text{HF})_2 \rightarrow \text{Na}\cdots\text{FH} + \text{HF}$, which is endoergic by roughly 0.07eV (the binding in $\text{Na}\cdots\text{FH}$ was calculated, *ab initio*, to be 1.8kcal/mole[110] whereas that for HF-HF has been measured as 2.97kcal/mole[111]). This process suggests another reason for the increased $\text{Na}\cdots\text{FH}$ concentration when He was used as the carrier gas. In the presence of a large excess of the inert gas, the collision energy will be larger for an expansion in He than in Ar. The final kinetic energy of the molecular beam following expansion from the pulsed source is given by $(5/2)kT$ [92]. This gives a velocity, v , for the halide molecules of $\sim 1700\text{m/s}$ in He and $\sim 500\text{m/s}$ in Ar. The relative collision energy of the molecular beams is $(1/2)\mu v^2$, where μ is the reduced mass. Therefore, for $\text{Na} + (\text{HF})_2$, the collision energy was $\sim 0.3\text{eV}$ when He was used as the carrier gas, compared with $\sim 0.1\text{eV}$ when Ar was used (the velocity of the Na beam was $\sim 1200\text{m/s}$ - see section 2.1). The much higher collision energy obtained using He as the carrier gas will significantly exceed the barrier for the endoergic reaction. These conditions are in contrast to those found optimum for formation of $\text{Na}\cdots\text{XCH}_3$ complexes which appeared to favour the lower collision energy when Ar was used as the carrier gas. A photoionization time-of-flight (TOF) mass spectrum for the formation of $\text{Na}\cdots\text{FH}$ complexes under the above conditions was shown in fig. 2.5.

Ab initio calculations on the $\text{Na}\cdots\text{FH}$ complex have been performed by Trenary et al.[42] at the self-consistent field (SCF) level, Paniagua and co-workers[75] who used the Hartree-Fock method, and Sevin et al.[76] who used SCF theory incorporating configuration interaction (CI). The

results of these indicated that the $\text{Na}\cdots\text{FH}$ complex is bound by only 1.5-2.5kcal/mol. The multi-reference configuration interaction calculation performed in this laboratory[110] gave a value of 1.8kcal/mol (see section 4.3). All the indications are, therefore, that the $\text{Na}\cdots\text{FH}$ complex is more weakly bound than $\text{Na}\cdots\text{FCH}_3$, for which *ab initio* calculations indicate a bond energy of ~4.8kcal/mol (see section 3.2). This is in accord with the experimental observation that a gas-mixture containing 2.5% FCH_3 or 2.5% FH in He gave a signal intensity 5× larger for $\text{Na}\cdots\text{FCH}_3$ than for $\text{Na}\cdots\text{FH}$. However, other factors, such as the collision-efficiency for complex formation may affect this comparison, as is evident from the foregoing discussion.

4.2 Photodepletion measurements

The depletion of the $\text{Na}\cdots\text{FH}$ ion signal by laser L_1 at a fixed wavelength, $\lambda_1 = 660\text{nm}$, was shown in fig 2.5. In this section the full action spectrum will be described; i.e. the effect of the tunable-excitation laser, L_1 , on the signal from the fixed-ionization laser, L_2 , at 248nm. The interpretation of the observed photodepletion will be discussed in section 4.3. In summary, the stable complex, $\text{Na}\cdots\text{FH}$ is excited to a reactive PES which would be traversed by the reaction $\text{Na}^* + \text{HF} \rightarrow \text{NaF} + \text{H}$. In the range of wavelengths studied in this work, the excitation will form the Transition State (TS) in configurations which correlate with the excited-state reaction of $\text{Na}^*(3p\ 2P)$ with FH .

In the action spectrum of fig. 4.2 the photodepletion cross-section, σ_{dep} , of the $\text{Na}\cdots\text{FH}$ complex is shown as a function of the excitation wavelength, λ_1 , at a fixed ionization wavelength, $\lambda_2 = 248\text{nm}$.

Figure 4.2: Action spectrum for the Na·FH complex, showing the depletion cross-section in Å² against the wavelength of laser L₁ in nm.

Photodepletion was measured to either side of the Na D-line (589nm), across an energy interval of $\sim 11000\text{cm}^{-1}$ or $\sim 1.3\text{eV}$. In addition, the spectral region from 440 to 550nm was also probed, but the depletion cross section was found to be negligible.

The action spectrum for $\text{Na}\cdots\text{FH}$ is dominated by a broad peak between 550 and 680nm which is characterized by a high cross-section. This dominant feature is asymmetric, with a relatively sharp onset at 680nm (1.82eV) and a gradual fall-off from its maximum at approx. 660nm (1.88eV) to zero in the region of 550nm (1.91eV). In addition there is a broad feature between 700 and about 830nm characterized by lower cross-section; $\sim 0.1\times$ that of the strong peak. Both peaks have reproducible structure, with a spacing suggestive of vibration. The structure in the strong photodepletion region is dominated by two sharp spikes, one at 652nm and one at 662nm. An explanation for the origin of these spikes will be given in section 4.3.

It is instructive to compare this action spectrum with that for the system $\text{Na}\cdots\text{FCH}_3$ described in section 3.3. The $\text{Na}\cdots\text{FCH}_3$ spectrum exhibited four photodepletion maxima, located at 415nm, 485nm, 590nm, and 685nm. The last two peaks, to either side of the Na D-line, were the most intense and were ascribed to excitation of the complex to TS configurations for the reaction of Na^* in the $3p\ ^2P$ excited state with FCH_3 . The two photodepletion peaks at higher excitation energies were believed to correlate with higher electronic states of Na^* , namely the $4s\ ^2S$ and $3d\ ^2D$ states. By contrast, for $\text{Na}\cdots\text{FH}$, both photodepletion peaks in the action spectrum were well to the red of the Na D-line with maxima at ~ 660 and 780nm , and the two peaks far to the blue of the D-line were

absent. The maximum photodepletion cross-section in the earlier Na $\cdot$ FCH₃ action spectrum was $\sim 0.7 \text{ \AA}^2$ at 590nm. A similar maximum cross-section was observed in the Na $\cdot$ FH action spectrum for the first peak to the red of the D-line.

Fine structure similar to that in fig. 4.2 was observed in this laboratory in the action spectra of Na $\cdot$ FPh[17] and Na $\cdot$ FCH₃ (see section 3.3). For both of these complexes a vibrational line spacing of $\sim 240 \text{ cm}^{-1}$ was observed for at least one progression. The structure was attributed in section 3.3 to vibrational levels within the excited state of the complex, in particular the Na $\cdot$ F stretching mode. However, clear vibrational progressions were observed in the photodepletion peaks for Na $\cdot$ FCH₃ and Na $\cdot$ FPh, unlike the structure in the Na $\cdot$ FH spectrum which appears more complex. It is likely either that a larger number of vibrational progressions are superimposed in the region of photodepletion studied for Na $\cdot$ FH, or that the vibrational lines are substantially perturbed.

The lifetime of the TS, [Na $\cdot$ FH] ‡ , can be estimated from measurements of the depletion intensity for Na $\cdot$ FH as a function of the power of the excitation laser, L_1 . In section 2.1 measurement of this depletion curve was described using an excitation wavelength of 660nm. A linear increase in the logarithm of the inverse of the depletion, $\ln(N_0/N)$ (where N , N_0 are the signal intensities in the presence and absence of L_1), was obtained with excitation-laser fluences, F_1 , up to $\sim 4 \text{ mJ/cm}^2$, above which saturation effects were observed. In section 2.3 the dependence of the depletion intensity on the excitation-laser fluence was given using a kinetic scheme that neglected laser-induced stimulated

emission. The resulting equation, $\ln(N_0/N) = \sigma F_1$, was noted only to be valid in the linear regime. In order to describe the effect of F_1 on the depletion signal at higher laser fluences, i.e. the full curve of fig. 2.7, the following kinetic scheme must be used.

where N is the concentration of the ground-state complex ($\text{Na}\cdot\text{FH}$), N^* is the concentration of TS species, $[\text{Na}^*\cdot\text{FH}]^\ddagger$, B is the Einstein coefficient, ρ_ν is the spectral density and $\tau^\ddagger$ is the lifetime of the TS. The quantity $B\rho_\nu$ can be equated to σI_1 where σ is the depletion cross-section for $\text{Na}\cdot\text{FH}$ measured in the linear regime and I_1 is the flux of the excitation laser in $\text{\AA}^{-2}\text{s}^{-1}$. The rate equations can be solved to give an explicit expression for the inverse depletion,

$$\ln (N_0/N) = \sigma F_1 -$$

$$\ln \left\{ 1 - \frac{\sigma F_1}{T} \left[1 - e^{-T} - \frac{1}{(\sigma F_1 / T + 1)e^{-\sigma F_1}} \left(1 - e^{-\sigma F_1} e^{-T} \right) \right] \right\} \quad (4.4)$$

where $T = t/\tau^\ddagger$ and t is the duration of the excitation-laser (L_1) pulse ($t \sim 20\text{ns}$). It has been assumed that the flux of L_1 is constant during the excitation pulse. The fluence, F_1 , is therefore equal to $I_1 t$.

The inverse depletion has been calculated as a function of the excitation-laser fluence at $\lambda_1 = 660\text{nm}$ using a series of values for the TS lifetime. The results are illustrated in fig. 4.3. An accurate description of

Figure 4.3: Power dependence of the depletion signal for the complex Na·FH at an excitation wavelength, λ_1 , of 660nm. The experimentally measured values are represented by circles. The dependence of $\ln(N_0/N)$ on F_1 has been calculated using a series of values for the TS lifetime, $\tau^\ddagger$.

the experimentally measured values was obtained when a lifetime of $\tau^{\ddagger}=3\times 10^{-9}\text{s}$ was used.

4.3 Discussion

Dynamics of the photoinduced reaction

Excitation of the $\text{Na}\cdots\text{FH}$ complex by laser L_1 can be described as being due to D-line excitation of the Na chromophore which is perturbed by the adjacent HF molecule. The fact that the major spectral features are shifted to the red from the D-line indicates that one or more excited states of the complex, correlating with the $\text{Na}(3p\ ^2P) + \text{HF}$ limit, are more strongly bound than the ground state. *Ab initio* computations[110] predict this to be the case (see below). Stronger binding in $\text{Na}^*\cdots\text{FH}$ can be explained simply as being related to the fact that $\text{Na}^*(3p\ ^2P)$ has a permanent quadrupole moment that can interact electrostatically with the dipole moment of HF[110] (the corresponding interaction potential has a R^{-4} dependence, where R is the distance between Na and the center of mass of HF). The ground-state ($3s\ ^2S$) of Na has no permanent multipole moments, so that only dispersion interactions and a dipole-induced-dipole interaction can stabilize the ground state of the $\text{Na}\cdots\text{FH}$ complex; both these forces have an interaction potential with a R^{-6} dependence.

Dissociation of $\text{Na}^*\cdots\text{FH}$ was described in ref. 76 to occur by 'hopping' to the ground state with resultant formation of the ionic-reaction product, Na^+F^- , or re-formation of the original $\text{Na} + \text{HF}$. For the special case of excitation to the blue of the Na D-line, the complex can also dissociate to form $\text{Na}^* + \text{HF}$. All these processes will be discussed in

terms of the electronic PESs of NaFH obtained by *ab initio* calculation[110].

The photoinduced reaction can be symbolized, starting with the ground-state complex, as

where the double-dagger represents the transition state, TS. The charge-transfer process can occur when the internuclear separations, $r_1(\text{Na-F})$, $r_2(\text{F-H})$ and $r_3(\text{Na-H})$, have reached a configuration of the TS in which the covalent and ionic wavefunctions are degenerate; it is symbolized by the bold arrow in eq. 4.5. This will be shown to be directly related to the surface-hop event described by Sevin et al.[76] which occurs in the TS for the $\text{Na}^* + \text{FH}$ reaction from the excited-state PES into the product channel, $\text{NaF} + \text{H}$, on the ground-state PES.

Near-degeneracy of the covalent and ionic wavefunctions is only expected to occur if r_2 is stretched[17,19,38]. Extension of r_2 (by $\sim 0.26\text{\AA}$; see ref. 51) is required to obtain a positive electron affinity for HF. At smaller r_2 , HF would repel electrons. However, at longer stretches of the H-F bond an electron can be transferred from Na^* to HF to form the ionic TS, $\text{Na}^+\cdots(\text{F}^-\text{--H})^-$. This is also evident in the recent *ab initio* results from this laboratory[110].

A simple rationale for the need to stretch F-H is that reaction on the ground-state PES is substantially endoergic and, consequently, has a late barrier located at an extended r_2 configuration. The crest of this barrier in the ground-state marks a region where the PESs for the ground

and first-excited states approach each other closely. As a consequence, the 'hop' from one PES to the other is most likely to occur at an extended r_2 . This hop can lead to reaction ($\rightarrow \text{Na}^+\text{F}^- + \text{H}$) if there is sufficient momentum along the r_2 coordinate. Otherwise the hop can lead to 'failed reaction' ($\rightarrow \text{Na} + \text{FH}$).

The adiabatic PESs of ground and electronically-excited states of $\text{Na}\cdots\text{FH}$ [110] were generated using the MRDCI variant of the multi-reference (MR) configuration-interaction (CI) method with single and double excitations[107,108]. The *ab initio* calculations performed in this laboratory are more extensive and give evidence of being more accurate than those published earlier. The full details of this calculation are described in ref. 110. Three excited states, with term symbols $2^2\text{A}'$, $1^2\text{A}''$ and $3^2\text{A}'$ (using the C_s point group to classify the electronic states of NaFH), arise from the interaction of Na in the $3p\ ^2\text{P}$ excited state with HF in the ground state.

Figure 4.4 gives the locations and energies of some characteristic points on the PES of the ground and excited states as a function of all three coordinates needed to describe the three-body system; $r_1(\text{Na-F})$, $r_2(\text{H-F})$, and the Na-F-H angle. These result from the fitting of MRDCI energies to an analytical representation described in ref. 110. The saddle point on the ground-state surface corresponds to the crest of the energy barrier on the minimal-energy path for the reaction and is located in the region of the avoided crossing between the PESs for the ground and first-excited state. The reaction $\text{Na} + \text{HF} \rightarrow \text{NaF} + \text{H}$ on the ground PES is 1.3eV endoergic and, as expected, the saddle point is located in the exit valley of the PES, with $r_2(\text{H-F})$ extended by $1.77 \times r_0(\text{H-F})$ and $r_1(\text{Na-F}) =$

Excited State ($3^2A'$, $1^2A''$)

potential energy
minimum

$$D(\text{Na}^*\cdots\text{FH}) = 0.33\text{eV}$$

Excited State ($2^2A'$)

potential energy
minimum

$$D(\text{Na}^*\cdots\text{FH}) = 0.532\text{eV}$$

Ground State ($1^2A'$)

(a) potential energy
minimum

$$D(\text{Na}\cdots\text{FH}) = 0.076\text{eV}$$

(b) saddle point

Figure 4.4: *Ab initio* MRDCI calculation[110] of the geometry and dissociation energy (D) of the most stable configurations on the ground-state ($1^2A'$) and excited-state ($2^2A'$) PES of NaFH. The geometry of the saddle-point on the ground-state PES is also shown. Equilibrium values for the bond lengths are $r_0(\text{Na-F})=1.926\text{\AA}$ and $r_0(\text{H-F})=0.917\text{\AA}$.

$r_0(\text{Na-F})$. In good agreement, with earlier semi empirical[74] and *ab initio*[76,77] calculations, the geometry at the saddle point is approx. 90° and at a height of 1.260eV (29kcal/mol) above the $\text{Na}+\text{HF}$ asymptote.

The saddle point on the lower PES was described above as corresponding approximately to the minimum energy-gap TS configuration for the reaction of Na atoms with HF. This is the configuration in which the ionic and covalent states of the system are most-nearly degenerate and is the probable location of the harpooning event in the reaction of alkali metals with halogen molecules (see section 1.3). The extended H-F bond length and the bent structure is consistent with stabilization of the ionic configuration, $\text{Na}^+\cdots\text{FH}^-$.

Figure 4.4 also lists well-depths for the $1^2\text{A}'$ and $2^2\text{A}'$ states of $\text{Na}\cdots\text{FH}$. The ground-state value can be compared to the well-depths determined by Düren and co-workers[69] for the non-reactive scattering of ground state Na from HF which were 17meV and 65meV. The former was likely to be due to scattering of HF from the H-end of the molecule. The latter can be compared to the well-depth of 76meV (1.8kcal/mol) for the $\text{Na}\cdots\text{FH}$ complex calculated here. The geometry of the most stable configuration of the ground and excited-state complex resulting from MRDCI calculations is given in fig. 4.4. It is possible that MRDCI slightly overestimates the binding in $\text{Na}\cdots\text{FH}$ due to basis-set superposition error, BSSE[110], which could not be accounted for at this level of theory. For example, a coupled-cluster theory with singles, doubles and perturbative triples (the so-called CCSD(T) approach) gives, after correcting for BSSE, 66.5eV[110]. The corresponding well-depths for the excited states, $\text{Na}^*\cdots\text{FH}$, are 532meV for the $2^2\text{A}'$ state and ~330meV for the $3^2\text{A}'$ and

$1^2A''$ states. These results are in accord with experimental estimates of 487 and 337 meV for the well depths determined by Düren and co-workers[69].

The collinear adiabatic PESs for the ground ($1^2A'$) and first-excited ($2^2A'$) states are illustrated in a 3D representation in fig.4.5. A schematic representation of the Transition State Spectroscopy dynamics proposed here is shown. A vertical transition of energy $h\nu_1$ from the region of the minimum on the ground-state PES accesses the excited-state in a region not far from an avoided crossing with the ground state. A possible pathway via surface-hopping to the ground-state product channel $\text{NaF} + \text{H}$ has been sketched onto the PES of fig. 4.5. Following the absorption of the visible light, $h\nu_1$, $\text{Na}^*\cdots\text{FH}$ is shown being formed with the $\text{Na}^*\cdots\text{F}$ bond extended. An oscillation in the upper PES leads, in the schematic, to a compressed $\text{Na}^*\cdots\text{FH}$ which hops down to the ground-state PES. The hop from the electronically-excited state occurs to the ground state in the region of the saddle point such that the electronic wavefunction is converted from largely covalent into largely ionic during the surface hop event. The hop is the discontinuous event identified as 'harpooning'. Since $\text{Na}^*\cdots\text{FH}$ is compressed on the lower PES, the product trajectory is shown oscillating across the exit valley as Na^+F^- is formed in a state of vibrational excitation.

Surface-hop events are nonadiabatic transitions which occur (with highest probability) at the seam of avoided crossing between two strongly coupled diabatic states underlying the adiabatic representation. The probability of the downward hop can be obtained, for example, from the Landau-Zener expression;

Figure 4.5. Adiabatic collinear PESs of the ground ($1^2A'$) and first-excited ($2^2A'$) states of NaFH generated by MRDCI calculation[110]. A schematic representation is given of the dynamics of photoexcitation of the Na··FH complex to the TS region of the excited state, followed by 'hopping' to the ground electronic-state to form ionic product, Na^+F^- .

$$\text{Pr (surface hop)} = \exp (-\pi^2\xi) \quad (4.6)$$

$$\text{where } \xi = \Delta E / (\hbar v |\Delta F|)$$

according to which the hopping probability decreases exponentially with the magnitude of the energy gap, ΔE , between the upper and lower adiabatic PES. However, the hopping probability also depends on the shape of the surfaces (ΔF is the difference in the gradient of the surfaces in the diabatic representation) and the time the system spends in the region of the avoided crossing (v is speed of the relative motion). Functional forms of the PESs for the ground and electronically-excited states have been generated in both the adiabatic and diabatic representations[110]. This will enable a trajectory-surface hopping (TSH) study of the evolution of the TS on the excited-state PES. Such a calculation is now in progress[128].

Another region of close approach of the ground and first excited-state with a much smaller energy gap (0.1 eV or less) is evident at greatly extended $r_1(\text{Na-F})$ and $r_2(\text{F-H})$. This affords the possibility of a second reactive pathway starting with substantial vibrational excitation in the F-H coordinate of $\text{Na}\cdots\text{FH}$, since the close approach occurs at extended r_2 as well as extended r_1 . The dynamics in this case would involve charge transfer as the $\text{Na}^*\cdots\text{FH}$ was about to separate into three atoms in the configuration $\text{Na}^*\cdots\text{F}\cdots\text{H}$. This would constitute the harpooning of a distant but slow-moving whale. The long-range Coulombic attraction between Na^+ and F^- subsequent to the harpooning event would provide the restoring force that brought NaF back together with extremely high vibrational energy. However, this reactive path is not accessible by a

Figure 4.6: Potential-energy curves for the ground and the electronically-excited states, along an approximate minimum-energy path for the collinear PES[110]. The equilibrium values for the Na-F and H-F bond lengths are indicated on their respective axes by downward pointing arrows at $r_1 = r_0(\text{Na-F}) = 1.926 \text{ \AA}$ and $r_2 = r_0(\text{H-F}) = 0.917 \text{ \AA}$. The minimum-energy gap between the ground and first-excited state lies between the two horizontal markers on the respective potential-energy curves.

vertical transition from stable configurations of the ground-state complex. It would require initial vibrational excitation of HF within the ground state complex prior to formation of the TS for the excited-state reaction.

Figure 4.6 gives a simplified view of the dynamics observed in the experiment, obtained by plotting the PE along the (approximate) minimum-energy path of the collinear ground-state PES. The smallest energy gap for the collinear geometry ($\sim 0.85\text{eV}$) is indicated by two horizontal markers. A hop downward at this $r_2(\text{H-F})$ internuclear separation would access the ground state ahead of the barrier crest. The system can, however, still surmount the barrier to reaction, to give $\text{NaF} + \text{H}$, provided that the necessary momentum is present. This momentum along r_2 (the exit valley) is obtainable most efficiently from vibrational motion in the HF bond under attack[37].

The observed depletion spectrum is plotted along the ordinate at the appropriate energy. The separation between the ground state PES ($1^2\text{A}'$) and the electronically-excited PESs ($2^2\text{A}'$, $1^2\text{A}''$, $3^2\text{A}'$), as computed *ab initio*, is in satisfactory agreement with the observed excitation energies for photodepletion. The peak at low energy (in the range $1.49\text{--}1.77\text{eV}$), in common with the major peak at $>1.8\text{eV}$, exhibits vibrational structure. In accord with fig. 4.6, they must constitute bound-to-bound transitions.

The presence of the two peaks in the experimental action spectrum, which is reproduced at the top left of fig. 4.6, can be understood in terms of the existence of three electronically-excited states

correlating with the $\text{Na}^*(3p\ ^2P) + \text{HF}$ asymptotic limit. The weak photodepletion peak below 1.8 eV arises from excitation of $\text{Na}\cdots\text{FH}$ to the first excited state, $2^2A'$. The peak above 1.8 eV in the observed action spectrum is seen to correspond to excitation of $\text{Na}\cdots\text{FH}$ to the higher excited states, namely $1^2A''$ and $3^2A'$. In this case depletion of the complex probably occurs via coupling of the $1^2A''$ and $3^2A'$ states with the $2^2A'$ state, followed by surface hopping from the $2^2A'$ state to the ground state, $1^2A'$. The fact that the tail of the strong photodepletion peak extends to an energy 0.1 eV above the dissociation limit for $\text{Na}^* + \text{FH}$ formation may be explained by an excitation of the bound-to-free variety. This should give rise to some D-line fluorescence (which we have not attempted to detect) through dissociation to yield $\text{Na}^* + \text{FH}$ which competes with reaction to give $\text{NaF} + \text{H}$.

Figure 4.6 gives the impression that the first electronic-excited state has a minimum which (though deep) is too high in energy to support the low-lying vibrational levels required to account for the observation of this low-energy ('red') peak in the photodepletion spectrum. It should be recalled, however, that fig. 4.6 gives the ground and excited PE curves along the collinear (minimum-energy) path, only. The complex is in fact considerably bent in the ground and electronic-excited states at its minimum energy conformations (see fig. 4.4).

Once transferred to the excited state, the complex, $[\text{Na}^*\cdots\text{FH}]^\ddagger$, can 'hop' back to the ground state through inter-state coupling. The electronically-excited states with A' symmetry shown in fig. 4.6 can couple with the ground state which also has A' symmetry. The 'harpooning' event corresponds to a nonadiabatic 'hop' from the excited

states of A' symmetry to the electronic-ground state. The outcome can be the ionic product $\text{Na}^+\text{F}^- + \text{H}$ or, if the energy and momentum along the reaction coordinate are insufficient, $\text{Na} + \text{FH}$.

The $2^2\text{A}'$ state is strongly coupled with the $1^2\text{A}'$ ground state in the region of close approach of the PES, in spite of a substantial energy gap ($\geq 0.85\text{eV}$) between the PESs. The $1^2\text{A}''$ excited state does not have the correct symmetry for nonadiabatic coupling with the ground state (even though it may couple with the $1^2\text{A}'$ state via Coriolis terms). The $3^2\text{A}'$ state, which does have the appropriate symmetry for nonadiabatic coupling, does not appear to approach the ground state sufficiently closely for strong coupling to occur. We believe, therefore, that the fastest pathway from the TS explored here, to the products is via the $2^2\text{A}'$ state.

A surface-hopping trajectory study on the NaClH system, performed by Yamashita and Morokuma[82], indicated that the lifetime of the system excited to the $2^2\text{A}'$ state is $\sim 1\text{ps}$. We anticipate a somewhat similar result for the NaFH system, indicating that the $2^2\text{A}'$ state is depopulated by surface-hopping long before significant fluorescence occurs. However, the excited state would need to be formed with sufficient internal energy such that the region of close approach between the ground and electronically-excited states can be reached. If the excited state is formed with insufficient internal energy then it may fail to deplete rapidly. A lifetime for the TS which was much longer than 1ps was observed at an excitation energy of 660nm . The lifetime of the TS formed for these photoinduced reactions will be shown in the next chapter to be strongly dependent on the excitation energy (for the related

example of $\text{Na}\cdots\text{FCH}_3$). Further studies will be required to elucidate the pathway for depletion following excitation to the higher excited states, $1^2\text{A}''$ and $3^2\text{A}'$.

Vibrational structure and intensity of the photodepletion peaks

An absorption spectrum for the $\text{Na}\cdots\text{FH}$ complex has been simulated by Topaler et al. in ref. 114 and is reproduced in fig. 4.7. The theoretical spectrum was generated using analytical fits to the *ab initio* potential energies for NaFH which were calculated in this laboratory using a large number of ground and excited-state configurations[110]. Once the adiabatic PESs for the $1^2\text{A}'$, $2^2\text{A}'$, $3^2\text{A}'$, and $1^2\text{A}''$ states were obtained in a functional form, the vibrational modes for the ground and excited-state complexes could be determined and Franck-Condon factors for vertical transitions between vibrational-quantum states could be obtained. The full details of the calculation are given in ref. 114. The theoretical spectrum was simulated at a number of different internal temperatures for the ground-state complex. At a temperature of 250K, a good qualitative agreement with the measured photodepletion spectrum was obtained (see fig. 4.7), although the absolute values of the cross-section showed a significant discrepancy. The features of the simulation which have enabled a better understanding of the measured photodepletion spectrum will be described below. The measured photodepletion spectrum would only be identical to the calculated absorption spectrum if *all* absorption led to depletion of the ground-state $\text{Na}\cdots\text{FH}$.

Figure 4.7: (a) Experimental photodepletion spectrum for the Na··FH complex as a function of the excitation energy (reproduced from fig. 4.2). (b) Calculated absorption spectrum for Na··FH at 250K[114]. (c) Calculated components of the total absorption cross section; $2^2A' \leftarrow 1^2A'$ (solid line), $1^2A'' \leftarrow 1^2A'$ (short-dashed line), and $3^2A' \leftarrow 1^2A'$ (long-dashed line)[114].

The depletion cross-section measured in the experimental spectrum (fig. 4.7(a)) was considerably less than the absorption cross-section calculated in the simulated spectrum (fig. 4.7(b)) at all excitation wavelengths. In addition, the relative intensity of the photodepletion peaks (which arise respectively from the transitions $2^2A' \leftarrow 1^2A'$ at $E_1 < 1.8\text{eV}$ and $1^2A'', 3^2A' \leftarrow 1^2A'$ at $E_1 > 1.8\text{eV}$) determined in the experimental spectrum show a significant discrepancy from the computed peaks in the absorption spectrum. This suggests that, for the observed excitation, there is a significant probability that the excited-state complex, $\text{Na}^* \cdots \text{FH}$ may fail to deplete. Furthermore, at low excitation energies, the probability that the excited-state complex fails to deplete is larger.

The integrated photoabsorption cross-section across the range of excitation energies which include the transitions; $1^2A'', 3^2A', 2^2A' \leftarrow 1^2A'$, was estimated in ref. 114 to be $\sim 1.1\text{\AA}^2\text{eV}$. This value should be compared to the integrated photodepletion cross-section determined from the experimental spectrum in fig. 4.7(a) which has a value $0.17\text{\AA}^2\text{eV}$ across the same range of excitation energies.

The computed integrated cross-section was calculated using equation 3.1 and assuming that the combined oscillator strength, $\sum_i f_i$, for $i = 1^2A'', 2^2A', 3^2A' \leftarrow 1^2A'$ is ~ 1 . This was justified, in ref. 114, since the oscillator strength for the corresponding atomic transition, $3p\ 2P \leftarrow 2s\ 2S$, in Na is approximately unity. However, in reality the Na atom may be sufficiently perturbed in the complex such that the oscillator strength for transitions to higher electronically-excited states make a significant contribution in the summation $\sum_i f_i = 1$ and, consequently,

the combined oscillator strength for the restricted set of electronic transitions observed in the photodepletion spectrum are <1 . Excitation to higher electronic states, such as $4^2A' \leftarrow 1^2A'$, was observed in the photodepletion spectrum for $\text{Na}\cdot\text{FCH}_3$ (see fig. 3.4, in which the $4^2A'$ excited state was labeled $4s\ ^2S$). The integrated cross-section for photodepletion from the $4^2A'$ state of $\text{Na}\cdot\text{FCH}_3$ (peak A in fig. 3.4) was $0.03\text{\AA}^2\text{eV}$ which should be compared with the value of $0.17\text{\AA}^2\text{eV}$ for photodepletion from the $1^2A''$, $2^2A'$ and $3^2A'$ states of $\text{Na}\cdot\text{FCH}_3$ (peaks B and C in fig. 3.4, in which the excited states were labeled $3p\ ^2P$). For the system $\text{Na}\cdot\text{FCH}_3$, it is expected that the oscillator strength for the excitation $4^2A' \leftarrow 1^2A'$ will make a significant contribution in the summation $\sum_i f_i = 1$. However, the contribution of this state for $\text{Na}\cdot\text{FH}$ would not account for the much larger discrepancy between the computed photoabsorption cross-section and the measured photodepletion cross-section.

A further mechanism which could in principle give rise to the discrepancy between the photoabsorption and the photodepletion cross-sections would involve laser-induced (stimulated) emission from the TS, which reforms the ground-state complex in a bound configuration. However, if this mechanism was operating then the dependence of the photodepletion signal on the fluence of the excitation laser, F_1 , would be non-linear. This was not observed in fig. 2.7 for the laser-fluences used in the measurements of the experimental depletion cross-section, so the stimulated emission pathway cannot be significant.

The origin of the two spikes in the strong photodepletion peak observed at the excitation wavelengths, λ_1 , of 652 and 662nm has been

determined. They were assigned in ref. 114 to vibrational transitions from the ground state ($1^2A'$) to the second ($3^2A'$) and third ($1^2A''$) electronically-excited states which overlap in the measured spectrum, giving rise to an enhanced cross-section at these wavelengths.

Vibrational progressions within the calculated components of the photoabsorption spectrum of $\text{Na}\cdots\text{FH}$ (fig 4.7(c)) can be identified. However, as a result of the presence of a large number of overlapping vibrational lines, the progressions become somewhat obscured in the combined calculated spectrum (fig 4.7(b)) and even more so in the experimental photodepletion spectrum (fig 4.7(a)). The experimental finding of somewhat indefinite vibrational progressions for $\text{Na}\cdots\text{FH}$ was in contrast to the action spectra for $\text{Na}\cdots\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$) in which clear vibrational progressions were visible.

The calculated fundamental frequencies for the ground and electronically-excited states of the complex are listed in table 4.1. The vibrational frequency for the F-H stretching mode was determined to be significantly reduced in the first-excited state, $2^2A'$ (from 3982cm^{-1} to 2889cm^{-1} [114]). The depth of the potential-energy well for the $2^2A'$ state is sufficient for bound excitation of the H-F vibrational mode within the complex, $\text{Na}^*\cdots\text{FH}$, according to the calculated frequency. This is not possible in any other of the calculated electronic states. Excitation of the HF vibrational mode in $[\text{Na}^*\cdots\text{FH}]^\ddagger$ was found, in ref. 114, to contribute to the photo-absorption spectrum at short wavelengths ($\lambda_1 < 600\text{nm}$ or $E_1 > 2.1$). In this portion of the spectrum the absorption cross-section was very small.

Table 4.1: Calculated frequencies for vibrational modes in ground¹ and excited² states of Na·FH[114].

Electronic state	1 ² A'	2 ² A'	1 ² A''	3 ² A'
<i>Na-F stretch</i> ¹ <i>/cm⁻¹</i>	115	252	269	269
<i>Na-F-H bend</i> <i>/cm⁻¹</i>	35	136	228	260
<i>F-H stretch</i> ² <i>/cm⁻¹</i>	3982	2889	4055	4055

¹the stretching frequency is 536cm⁻¹ in the NaF molecule[98].

²the stretching frequency is 4138cm⁻¹ in the HF molecule[98].

The knowledge that has recently been obtained of the adiabatic PESs will enable a detailed study by the surface-hopping trajectory method, leading to a better understanding of the dynamics which follow excitation of the $\text{Na}\cdots\text{FH}$ complex. A study of this sort is underway[128].

4.4 Summary

The van der Waals complex $\text{Na}\cdots\text{FH}$ has been observed for the first time in a crossed-beam apparatus. Reaction of the complex was found to be initiated by photoexcitation across a broad spectral region from 550-840nm, to either side of the sodium D-line (589nm). The photodepletion spectrum consisted of two broad peaks, one intense peak with its maximum shifted approx. 70nm to the red of the D-line and a weaker peak with its maximum shifted approx. 200nm to the red of the D-line. The substantial red-shift is indicative of stronger binding in the electronically-excited states than in the ground state. The reactive excited state of the complex (the 'transition state') lasted for a sufficient length of time to yield vibrational structure.

Both peaks observed in the action spectrum arose from photoexcitation to a potential-energy surface (PES) which would be traversed by a collision between Na^* in its $3p\ ^2P$ state with an HF molecule. Accurate *ab initio* calculations have indicated the presence of three electronically-excited states of $\text{Na}^*\cdots\text{FH}$ which correlate with the $\text{Na}^*(3p\ ^2P) + \text{HF}$ asymptotic limit. These can best be labeled according to the symmetry of the excited state described by the C_s point group; $2^2A'$, $3^2A'$ and $1^2A''$. The energy splitting between the electronically-excited states of the complex, $\text{Na}\cdots\text{FH}$, derived from the observed excitation

energies in the measured depletion of $\text{Na}\cdots\text{FH}$ agree with the computed separation between the PES for the ground ($1^2\text{A}'$) and excited ($2^2\text{A}'$, $3^2\text{A}'$, $1^2\text{A}''$) states[110]. The weak peak observed extending from 830 to 700nm corresponds to excitation to the lowest electronically-excited state of $\text{Na}^*\cdots\text{FH}$, $2^2\text{A}'$. The strong peak, measured from 700 to 550nm, is believed to arise from a photoinduced transition to the two higher excited-state PESs.

The adiabatic PESs obtained by *ab initio* calculation in ref. 110 indicated clearly that photoexcitation of $\text{Na}\cdots\text{FH}$ occurred to configurations intermediate between the reagent, $\text{Na}^* + \text{FH}$, and product, $\text{NaF} + \text{H}$, on the excited-state. This supported the description of the photodepletion experiment as a Transition State Spectroscopy (TSS) study of the harpooning reaction in which the TS was formed in different starting configurations as the excitation wavelengths was varied.

The region of closest-approach of the first-excited PES and the ground-state PES is situated adjacent to the barrier crest on the lower PES. The lower PES, as expected for a substantially endoergic reaction, $\text{Na} + \text{HF} \rightarrow \text{NaF} + \text{H}$ $\Delta H^\circ = 1.3\text{eV}$, has a 'late' barrier-crest, corresponding to stretching of the H-F bond under attack. Accordingly the region of closest approach of the upper and lower PES would be expected to occur at extended $r_2(\text{H-F})$. This finding had been anticipated earlier[17,19] on the grounds that extension of $r_2(\text{H-F})$ by stabilizing HF^- favors the charge-transfer needed for harpooning to occur. It is confirmed by *ab initio* calculations performed in this laboratory[110].

Chapter 5 Transition State ionization of $\text{Na}\cdot\cdot\text{FCH}_3$

A different approach to Transition State Spectroscopy of the reaction $\text{Na}^* + \text{XR} \rightarrow \text{NaX} + \text{R}$ would be to excite the ground-state complex, $\text{Na}\cdot\cdot\text{XR}$, to the potential-energy surface (PES) for the excited-state reaction and then to probe the decay of the Transition State (TS), $[\text{Na}^*\cdot\cdot\text{XR}]^\ddagger$, as it dissociates into products. This type of experiment has proved informative when performed in the time-domain. For example, the lifetime of the TS for the reaction $(\text{NaI})^* \rightarrow \text{Na} + \text{I}$ has been measured directly by Zewail and co-workers[30]. The same type of information has been obtained from experiments performed in the frequency-domain. The lifetime of the TS, $[\text{FNa}_2]^\ddagger$, for the reaction $\text{F} + \text{Na}_2 \rightarrow \text{NaF} + \text{Na}^*(3p\ 2P)$ was found by measuring the intensity of the emission to either side of the Na D-line[115]. The integrated intensity of the wing emission which arose from the TS, $[\text{FNa}_2]^\ddagger$, was found to be $\sim 10^{-4} \times$ that for D-line emission originating in the product, $\text{Na}^*(3p\ 2P)$. The lifetime of the $3p\ 2P$ state of Na is $\sim 10^{-8}$ s, giving a lifetime for the TS of $\sim 10^{-12}$ s.

In this chapter, an experiment performed in the frequency-domain will be described which has provided an order-of-magnitude estimate of the lifetime for the TS in the harpooning reaction. Monochromatic excitation of $\text{Na}\cdot\cdot\text{XR}$ complexes, using a tunable-dye laser, forms the TS, $[\text{Na}^*\cdot\cdot\text{XR}]^\ddagger$, for the excited-state reaction, $\text{Na}^* + \text{XR} \rightarrow \text{Products}$, in a selected-starting configuration. If it is assumed that all configurations of the TS lead to depletion of $\text{Na}\cdot\cdot\text{XR}$, then the kinetic scheme is given by

where $TS(\lambda_1)$ represents the starting configuration of the TS for an excitation wavelength to the TS of λ_1 . The depletion cross-section for $Na\cdots XR$, σ_{dep} , and the lifetime, $\tau^\ddagger$, will both depend on the initial configuration of the TS and, therefore, on the excitation wavelength, λ_1 . The concentration of TS species, $[TS]$, during the excitation-laser pulse is given implicitly by the rate equations,

$$\frac{d}{dt} [Na\cdots XR] = -\sigma_{dep} I_1 \times [Na\cdots XR] \quad (5.2)$$

$$\frac{d}{dt} [TS] = \sigma_{dep} I_1 \times [Na\cdots XR] - \frac{1}{\tau^\ddagger} \times [TS] \quad (5.3)$$

where $[Na\cdots XR]$ is the concentration of the ground-state complex and I_1 is the flux in photons/(cm²s) for the excitation laser, L_1 . These equations can be solved to give an explicit expression for $[TS]$,

$$[TS] = [Na\cdots XR]_0 \times \frac{\sigma_{dep} I_1}{1 / \tau^\ddagger - \sigma_{dep} I_1} \times \left(e^{-\sigma_{dep} I_1 t} - e^{-t/\tau^\ddagger} \right) \quad (5.4)$$

where $[Na\cdots XR]_0$ is the initial concentration of the ground-state complex.

The average concentration of TS species present during the excitation-laser pulse, can be estimated via ionization of $[Na^+\cdots XR]^\ddagger$ using a second laser, L_2 , which is fired at the same time as L_1 . The wavelength of L_2 must be sufficient long in order that ionization from the ground-state complex will not be observed. The ion signal for $Na^+\cdots XR$ can then be measured by time-of-flight (TOF) mass spectrometry. According to eq. 5.4, the measured concentration of TS species will be directly related to the lifetime for the TS.

Measurements of ionization from the TS; $[\text{Na}^*\cdots\text{FCH}_3]^\ddagger$, for the reaction $\text{Na}^* + \text{FCH}_3 \rightarrow \text{NaF} + \text{CH}_3$ have been made as part of the present experimental study. An action spectrum of the ion-signal intensity was recorded as a function of the excitation wavelength, λ_1 , used to initiate reaction in the $\text{Na}\cdots\text{FCH}_3$ complex. The TS ionization spectrum, which depends on the lifetime of the TS, will be compared with the depletion spectrum described in section 3.3 which is dependent only on the rate of removal of $\text{Na}\cdots\text{FCH}_3$ and not on the steady-state concentration, $[\text{TS}]$.

5.1 Experiment

The experiment was identical in principle to that described in chapter 2 except for the wavelength of the ionization laser, L_2 , and the time delay between L_2 and the excitation laser, L_1 . In brief, a wavelength of 343nm was used for L_2 , and the laser pulses were coincident in time (rather than L_2 being substantially delayed following L_1).

In order to record the photodepletion spectrum reported in section 3.3 above, the concentration of $\text{Na}\cdots\text{FCH}_3$ was monitored by an ionization laser at $\lambda_2=248\text{nm}$ (corresponding to an energy, $E_2=5.0\text{eV}$) capable of ionizing the complex in the electronic-ground state. By contrast, to obtain the TS ionization spectrum this laser was set at a longer wavelength of $\lambda_2=343\text{nm}$ (corresponding to an energy, $E_2=3.6\text{eV}$) which was insufficient to ionize the complex in the ground state but sufficient to ionize from an electronically-excited state, i.e. $[\text{Na}^*\cdots\text{FCH}_3]^\ddagger$.

The excitation and ionization wavelengths were provided by dye lasers (Lambda Physik LPD3000). The wavelength region used for the excitation laser, L_1 was covered by the laser dyes: LDS 698 (DMSO

solvent, 760-670nm), DCM (DMSO solvent, 690-632nm), Rhodamine 610 (Methanol solvent, 644-588nm), Coumarin 153 (Methanol solvent, 600-522nm). The ionization laser, L_2 , was operated using p-terphenyl dye (Exciton) in dioxane solvent.

In order to probe the TS for the photoinduced reaction it was necessary that the excitation-laser pulse, L_1 , and the ionization-laser pulse, L_2 , arrive in the interaction region at the same time. The signal from the ionization laser would then provide a measure of the average concentration of TS species which existed during the excitation-laser pulse. A single pump source was used to operate both dye lasers in order to produce synchronized laser pulses for L_1 and L_2 . Using a single source as the pump avoided the time-jitter which can arise from two separately triggered lasers. The output from an excimer laser (Lambda Physik LPX200) was split equally by a dielectric mirror, providing each dye laser with a pump beam. The excimer-pump laser was operated with a Xe/HCl/Ne gas mixture which gives rise to emission from XeCl^* at 308nm. The arrangement is illustrated in fig. 5.1.

The ionization laser, L_1 , was counterpropagated with the excitation laser, L_2 . The laser path of L_2 had to be extended by reflecting the beam back and forth in order to obtain identical path lengths and hence time-coincidence for each laser to the center of the extraction region of the TOF mass spectrometer. Both lasers were mildly focused and aligned such that the geometric cross-section for the ionization laser was overlapped with that for the excitation laser. The pulse width of both lasers was the same (~20ns).

Figure 5.1: Top view of the vacuum system and laser arrangement.

The TS ionization signal was measured across a broad wavelength range from 740-540nm using a step size of 0.25nm. The ion-signal intensity was normalised, following each laser pulse, to the excitation-laser fluence, F_1 , which was typically allowed to vary between ~ 1.5 to 2.5 mJ/cm^2 (i.e. below the saturation limit for photoexcitation noted in section 2.1). The wavelength of L_2 was fixed at $\lambda_2=343\text{nm}$ and the ionization-laser fluence, F_2 , was $\sim 20\text{mJ/cm}^2$. The TS ionization signal was found to be linearly dependent on the fluence of L_2 about this value. Fluctuations in the power of L_2 and the concentration of complexes formed in the crossed-molecular beams were corrected for by measuring the ion signal at a fixed wavelength for L_1 at regular intervals during the experiment. The intensity of TS ionization in the action spectrum was normalised according to this measured signal.

5.2 Results

The TS ionization signal was measured across a range of excitation wavelengths, λ_1 , from 740 to 540 nm, which correspond to excitation energies, E_1 , of 1.68 to 2.30 eV. This range included the region in which peaks B and C were observed in the photodepletion spectrum for $\text{Na}\cdots\text{FCH}_3$ (see fig 3.4). In this section, the TS ionization spectrum will be compared with the photodepletion spectrum, and a qualitative description of the evolution of the TS for the reaction $\text{Na}^* + \text{FCH}_3 \rightarrow \text{NaF} + \text{CH}_3$ will be given, based on these experimental findings.

The action spectra are illustrated in fig. 5.2 where the photodepletion spectrum (above) is a measure of the fraction of complexes which are removed by excitation at a particular wavelength,

Figure 5.2: (a) The photodepletion spectrum and (b) the TS ionization spectrum for the $\text{Na} \cdots \text{FCH}_3$ complex, as a function of the excitation energy, E_1 . The energy of the ionization photon, E_2 , was (a) 5.0 eV (248 nm) and (b) 3.6 eV (343 nm).

and the magnitude of the signal in the TS ionization spectrum (below) depends directly upon the lifetime of the TS, $[\text{Na}\cdots\text{FCH}_3]^\ddagger$. In these spectra, the excitation is given as energy in eV, instead of as the wavelength of laser L_1 . This region of the photodepletion spectrum, which includes peaks B and C, has been reproduced from fig. 3.4. A strong signal intensity appeared in the TS ionization spectrum only at the short wavelength side of peak C. At longer wavelengths and in the region of peak B, the signal intensity in the TS ionization spectrum was negligible. There was a small section of the TS ionization spectrum ($E_1 = 2.07\text{--}2.11\text{ eV}$) which was not recorded since across this region (which included the D-line at 2.11 eV), a strong ion signal was obtained for Na that saturated the detector, preventing measurement of the ion signal for $\text{Na}\cdots\text{FCH}_3$.

The spectra, fig. 5.2 (a) and (b) respectively, differed in each of the excitation-wavelength regions labeled (i), (ii) and (iii). The differences gave direct evidence of differing lifetimes for the TS. Thus in region (i) the TS, $[\text{Na}^*\cdots\text{FCH}_3]^\ddagger$, was formed with such low concentration and internal energy that it failed to deplete $\text{Na}\cdots\text{FCH}_3$ (fig. 5.2 (a); region (i)) but lingered in TS configurations for long enough to be ionized (fig. 5.2 (b); region (i)). In region (ii) with increasing excitation energy there was increasing depletion of $\text{Na}\cdots\text{FCH}_3$ (fig. 5.2 (a); region (ii)) but the concentration of the TS (fig. 5.2 (b); region (ii)) remained relatively constant, probably because the excitation efficiency was increasing sufficiently rapidly with excitation energy to offset the increasing reaction probability. Finally, in region (iii) the high energy to which $\text{Na}\cdots\text{FCH}_3$ was excited in the TS ensured efficient reaction (large

depletion in fig. 5.2 (a), region (iii)) despite the fact that the concentration of TS species was minimal (see fig. 5.2 (b); region (iii)). A quantitative estimate for the order-of-magnitude of the TS lifetime in regions (i), (ii), and (iii) will be made in section 5.3.

5.3 Discussion

The ion signal observed with an ionization energy of 3.6eV in fig. 5.2(b) cannot be due to ionization of ground-state $\text{Na}\cdots\text{FCH}_3$, but must be due to photoionization of the TS, $[\text{Na}^+\cdots\text{FCH}_3]^\ddagger$,

The kinetic equations will be solved to determine the magnitude of the ion signal, $N[\text{Na}^+\cdots\text{FCH}_3]$, measured in the TOF spectrum as a function of the lifetime for the TS.

The rate of formation of the complex ion is given by

$$\frac{d}{dt} [\text{Na}^+\cdots\text{FCH}_3] = \sigma_{\text{ion}} I_2 \times [\text{TS}] \quad (5.8)$$

and the ion signal is found by integration across the pulse width for the laser interaction (i.e. $0 < t < T$),

$$N[\text{Na}^+\cdots\text{FCH}_3] = \sigma_{\text{ion}} I_2 \times \int_0^T [\text{TS}] dt \quad (5.9)$$

It is assumed that the photon flux, I_2 , remains constant across the period of the laser pulse.

The TS is formed in selective-starting configurations by monochromatic excitation of the ground-state complex,

$\text{Na}\cdot\cdot\text{FCH}_3 + h\nu_1 \rightarrow \text{TS}(v_1)$, and decays to give products with a lifetime, $\tau^\ddagger$, which is dependent on the starting configuration. The excitation laser was synchronized with the ionization laser and both laser-pulse widths were approximately the same. The rate expression for the concentration of TS species in the presence of the ionization laser is given by

$$\frac{d}{dt} [\text{TS}] = \sigma_{\text{dep}} I_1 \times [\text{Na}\cdot\cdot\text{FCH}_3] - \frac{1}{\tau^\ddagger} \times [\text{TS}] - \sigma_{\text{ion}} I_2 \times [\text{TS}] \quad (5.10)$$

The concentration of ground-state complexes, $[\text{Na}\cdot\cdot\text{FCH}_3]$, is given by eq. 5.3, and the time-dependent concentration of TS species can be determined,

$$[\text{TS}] = [\text{Na}\cdot\cdot\text{FCH}_3]_0 \times \frac{\sigma_{\text{dep}} I_1}{1 / \tau + \sigma_{\text{ion}} I_2 - \sigma_{\text{dep}} I_1} \times \left(e^{-\sigma_{\text{dep}} I_1 t} - e^{-(1/\tau + \sigma_{\text{ion}} I_2)t} \right) \quad (5.11)$$

Again, it has been assumed that the photon flux for each laser, I_1 and I_2 , remains constant across the width of the laser pulses.

The ion signal can be calculated by inserting eq. 5.11, into eq. 5.9.

$$N[\text{Na}^+\cdot\text{FCH}_3] = [\text{Na}\cdot\cdot\text{FCH}_3]_0 \times \frac{\sigma_{\text{dep}} I_1 \times \sigma_{\text{ion}} I_2}{1 / \tau + \sigma_{\text{ion}} I_2 - \sigma_{\text{dep}} I_1} \times \left(\frac{1}{\sigma_{\text{dep}} I_1} \times \left\{ 1 - e^{-\sigma_{\text{dep}} I_1 T} \right\} - \frac{1}{1 / \tau + \sigma_{\text{ion}} I_2} \times \left\{ 1 - e^{-T / \tau} e^{-\sigma_{\text{ion}} I_2 T} \right\} \right) \quad (5.12)$$

where T is the width of the pair of laser pulses which corresponds to a time interval of $\sim 20\text{ns}$.

The rate constant for photoexcitation, $\sigma_{\text{dep}} I_1$, of the complex, $\text{Na}\cdot\cdot\text{FCH}_3$, was of the order of 10^7 s^{-1} . This was calculated using typical

values for the cross-section for photodepletion of $\text{Na}^+\cdot\text{FCH}_3$, $\sigma_{\text{dep}} \sim 0.25 \text{ \AA}^2$, and the excitation-laser fluence, $F_1 \sim 2 \text{ mJ}/(\text{cm}^2 \text{ pulse})$ (or $0.8 \text{ photons}/(\text{ \AA}^2 \text{ pulse})$). The value of F_1 corresponds to a photon flux, I_1 , of $\sim 4 \times 10^7 \text{ photons}/(\text{ \AA}^2 \text{ s})$.

The rate constant for ionization, $\sigma_{\text{ion}} I_2$, of the TS, $[\text{Na}^+\cdot\text{FCH}_3]^\ddagger$, was of the order of 10^8 s^{-1} . The photoionization cross section for Ar atoms has been measured as $\sim 50 \text{ Mbarns}$ (or $\sim 0.5 \text{ \AA}^2$) at photon energies which are just above the ionization limit [116]. Since ionization of Ar and $\text{Na}^+\cdot\text{FCH}_3$ both involve the removal of a 3p electron, it has been assumed that the photoionization cross section of $[\text{Na}^+\cdot\text{FCH}_3]^\ddagger$ has a similar value. The ionization-laser fluence, F_2 , was $\sim 20 \text{ mJ}/(\text{cm}^2 \text{ pulse})$ (or $4 \text{ photons}/(\text{ \AA}^2 \text{ pulse})$). The value of F_2 corresponds to a photon flux, I_2 , of $\sim 2 \times 10^8 \text{ photons}/(\text{ \AA}^2 \text{ s})$.

The estimated ion signal calculated using eq. 5.12 is shown in fig. 5.3 as a function of the lifetime of the TS, $\tau^\ddagger$. In order to observe significant ionization from the TS, the lifetime of the TS must be sufficiently long so that the concentration, $[\text{TS}]$, is detectable. A rapid decline in the ion signal is observed in fig. 5.3 when the lifetime of the TS is $< 10^{-8} \text{ s}$. This is likely to correspond to the sharp decline observed in the TS ionization signal when the excitation energy was increased above 1.8 eV. This leads us to estimate TS lifetimes, $\tau^\ddagger$, for regions (i), (ii) and (iii) in fig. 5.3 of $> 10^{-8}$, $\sim 10^{-8}$ and $< 10^{-8} \text{ s}$, respectively.

The calculated expression for the ion signal in eq. 5.12. can be simplified in the limit of long and short TS lifetimes, $\tau^\ddagger$. For a long lifetime (i.e. $\tau^\ddagger \gg 1/\sigma_{\text{ion}} I_2 \approx 10^{-8} \text{ s}$), the complex-ion signal is relatively

Figure 5.3: Estimated dependence of the ion signal from direct photoionization of the TS, $[\text{Na}^+ \cdots \text{FCH}_3]^\ddagger$, on the lifetime of the TS, using a typical photoionization cross-section of $\sim 0.5 \text{\AA}^2$.

constant,

$$N_{\{\text{Na}^+\cdots\text{FCH}_3\}} \approx 0.1 \times [\text{Na}\cdots\text{FCH}_3]_0 \quad (5.13)$$

and for a short lifetime (i.e. $\tau^\ddagger \ll 1/\sigma_{\text{ion}}I_2 \approx 10^{-8}\text{s}$), the complex-ion signal is proportional to the TS lifetime,

$$N_{\{\text{Na}^+\cdots\text{FCH}_3\}} \approx [\text{Na}\cdots\text{FCH}_3]_0 \times 10^7\text{s} \times \tau^\ddagger(/s) \quad (5.14)$$

The decline in the TS lifetime between regions (ii) and (iii) would be consistent with an energy barrier on the excited-state PES between the potential-energy minimum for $\text{Na}^+\cdots\text{FCH}_3$ and the region where surface-hopping that removes $\text{Na}^+\cdots\text{FCH}_3$ can occur.

The origin of the observed depletion in region (ii) is likely to be laser-induced stimulated emission from $[\text{Na}^+\cdots\text{FCH}_3]^\ddagger$ to dissociative configurations on the ground-state PES. In region (i), in which depletion was not observed, a bound configuration of $\text{Na}\cdots\text{FH}$ on the ground-state PES must be formed.

In region (iii), the TS may be formed with sufficient internal energy that it can access configurations on the excited-state PES from which surface hopping (see section 4.3) can occur to the ground-state product channels, $\text{NaF}+\text{CH}_3$ and $\text{Na}+\text{FCH}_3$. This would explain the fact that the excited state decays at a much faster rate in this region.

5.4 Summary

A Transition State Spectroscopy of the reaction $\text{Na}^++\text{FCH}_3 \rightarrow \text{NaF}+\text{CH}_3$ was performed in which the ground-state complex, $\text{Na}\cdots\text{FCH}_3$, was excited to the potential-energy surface (PES) for the excited-state

reaction and then the concentration and hence the approximate lifetime of the Transition State (TS), $[\text{Na}^+\cdots\text{FCH}_3]^\ddagger$, was obtained by direct photoionization from the TS.

The TS ionization signal was measured as a function of the excitation energy from 1.68 to 2.30 eV ($\lambda_1 = 740\text{-}540\text{nm}$). The action spectrum for photodepletion of $\text{Na}\cdots\text{FCH}_3$ in this wavelength range indicated peaks which differed from those in the TS ionization spectrum. The TS ionization signal depends on the concentration of TS species, and therefore the lifetime of the TS. On the basis of the ionization-laser fluence required in order to observe significant ionization from the TS, the lifetime of the TS must be $\sim 10^{-8}\text{s}$ in the spectral regions where TS ionization was observed. Comparison of this spectrum with that for photodepletion of $\text{Na}\cdots\text{FCH}_3$ led to the identification of three spectral regions in which the lifetime for dissociation of the TS was very different. At low excitation energies in the photodepletion spectrum, the lifetime of the TS was $>10^{-8}\text{s}$. As the excitation energy was increased, regions of TS configurations were accessed for which the lifetime of the TS was $\sim 10^{-8}\text{s}$. At the highest excitation energies in the range covered by the 'photodepletion spectrum' ($\text{Na}\cdots\text{FCH}_3 + h\nu_1 \rightarrow \text{TS} \rightarrow \text{products}$) the lifetime of the TS was found to be very short, $<10^{-8}\text{s}$.

Chapter 6 Conclusions and future directions

This thesis describes results of an experimental study of the photoinduced charge-transfer dissociation reaction in the van der Waals complexes, $\text{Na}\cdots\text{XR}$ ($\text{X} = \text{F}, \text{Cl}, \text{Br}$, $\text{R} = \text{CH}_3$; and $\text{X} = \text{F}$, $\text{R} = \text{H}$). For the case of $\text{Na}\cdots\text{FH}$ these results can be related to an extensive *ab initio* study of the same process performed (by others) in this laboratory. Photodepletion of these complexes is best understood as having occurred through excitation of the sodium chromophore followed by charge-transfer to the halide molecule, e.g. $\text{Na}\cdots\text{XR} + h\nu_1 \rightarrow [\text{Na}^*\cdots\text{XR}]^\ddagger \rightarrow [\text{Na}^+\cdots\text{XR}^-]^\ddagger \rightarrow \text{NaX} + \text{R}$, or $\text{Na} + \text{XR}$.

6.1 Conclusion

Photoinduced reaction in $\text{Na}\cdots\text{XCH}_3$ ($\text{X} = \text{F}, \text{Cl}, \text{Br}$) complexes was found to occur across a broad range of excitation wavelengths from $\sim 750\text{nm}$ to $\sim 390\text{nm}$, corresponding to an energy interval of $\sim 1.5\text{eV}$. The depletion cross section for $\text{Na}\cdots\text{XCH}_3$ was measured as a function of the excitation wavelength. These spectra consisted of up to four broad peaks, which were interpreted as arising from the photoinitiated reaction of XCH_3 with successively higher electronically-excited states of $\text{Na}^*(i)$, i.e. $i = 3p\ ^2P$, $4s\ ^2S$, $3d\ ^2D$, as the excitation energy was increased.

The $3p\ ^2P$ excited state of Na was found to be split in the environment of the complex giving rise to a pair of distinct photodepletion peaks. The energy spacing between the peaks was found to be 0.28eV for $\text{Na}\cdots\text{FCH}_3$, 0.47eV for $\text{Na}\cdots\text{ClCH}_3$ and 0.25eV for $\text{Na}\cdots\text{BrCH}_3$. For the case of $\text{Na}\cdots\text{ClCH}_3$ and $\text{Na}\cdots\text{BrCH}_3$, the two peaks were observed to either side of the Na D-line which corresponds to the $3p$

$2P \leftarrow 3s\ 2S$ transition in atomic Na. However, for $\text{Na}\cdots\text{FCH}_3$, the onset for depletion from both states was observed to the red of the D-line.

The location of the peaks in the photodepletion spectra can be understood in terms of the electronic states of the complex. These in turn arise from the atomic states of Na^* . The $3p\ 2P$ state is split into components which correspond to different orientations of the occupied $3p$ orbital relative to the closed-shell XCH_3 molecule. The photodepletion peak to the red of the D-line can be ascribed to reaction following excitation of an electron into a $3p$ orbital with a perpendicular orientation relative to the internuclear axis (labeled as the $3p\ 2P_{x,y}$ state). The location of this peak in the experimental photodepletion spectrum indicates that (in accord with theoretical calculations[95]) binding in the excited state is much stronger than binding in the ground-state complex. However, the $3p\ 2P_z$ state, in which the occupied $3p$ orbital is oriented along the internuclear axis is less strongly bound for $\text{Na}^*\cdots\text{ClCH}_3$ and $\text{Na}^*\cdots\text{BrCH}_3$ along the $\text{Na}\cdots\text{X}$ coordinate. This was reflected in the experimental action spectra for these complexes in which the photodepletion peak corresponding to this state was located to the blue of the Na D-line. However, the $3p\ 2P_z$ state was found to be weakly bound in $\text{Na}^*\cdots\text{FCH}_3$ and, consequently, photodepletion from this state was measured to the red of the D-line in the experimental action spectrum.

The *ab initio* calculation [96] indicated that the minimum-energy geometries for the $3s\ 2S$ (ground) and $3p\ 2P_{x,y}$ (first-excited) states of the complex are bent. This should give rise to a splitting within the $3p\ 2P_{x,y}$ state of the complex as a result of the CH_3 umbrella head. This was not

resolved in the experimental spectra although two pieces of indirect evidence did exist: (1) the photodepletion peak for the $3p\ 2P_{x,y}$ state was noted for being much broader than that for the $3p\ 2P_z$ state, particularly in the $\text{Na}\cdot\text{ClCH}_3$ action spectrum; (2) the photodepletion peak for the $3p\ 2P_{x,y}$ state of $\text{Na}\cdot\text{FCH}_3$ and $\text{Na}\cdot\text{BrCH}_3$ were both found to contain two sets of vibrational progressions which may each originate from different electronically-excited states.

The measured probability with which photodepletion occurred on each excited PES was found to increase with electronic excitation in Na (i.e. for $\text{Na}\cdot\text{FCH}_3$ from ~9% for $3p\ 2P_{x,y}$, to ~30% for $3p\ 2P_z$, to ~40% for $4s\ 2S$, and to ~100% for $3d\ 2D$) and, also, down the series of methyl-halide molecules, in the order $\text{Na}\cdot\text{FCH}_3 < \text{Na}\cdot\text{ClCH}_3 < \text{Na}\cdot\text{BrCH}_3$, for a particular electronically-excited state.

Vibrational structure within the photodepletion peaks was associated with stretching and bending modes in the TS. For $\text{Na}\cdot\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$), the vibrational structure within the lowest-energy photodepletion peak arising from the $3p\ 2P_{x,y}$ excited state was associated with the $\text{Na}\cdot\text{X}$ stretching mode. The structure observed in the second photodepletion peak arising from the $3p\ 2P_z$ excited state was associated with the $\text{Na}\cdot\text{X-CH}_3$ bending mode.

Photodepletion of the complex $\text{Na}\cdot\text{FH}$ was recorded across a similar energy interval to that which gave photodepletion for $\text{Na}\cdot\text{XCH}_3$. However, only two peaks were observed in the action spectrum. These arose from excitation to a PES which would be traversed by the reaction of Na^* in the $3p\ 2P$ state with the HF molecule. Both these peaks were

located to the red of the Na D-line transition indicating that binding in the excited-state is stronger than in the ground-state complex

Dissociation of the complex was believed to occur via a surface hopping mechanism from the excited-state PES into a product channel on the ground-state PES. The surface hop can occur in the region in which the ground and excited state approach each other closely.

A second action spectrum was obtained for the $\text{Na}\cdots\text{FCH}_3$ complex in which the TS, $[\text{Na}^*\cdots\text{FCH}_3]^\ddagger$, was probed by direct ionization. The cross-section for ionization of the TS permitted the lifetime of the TS to be estimated. It was found that at low excitation energies (1.67-1.70eV) the lifetime of the TS exceeded 10^{-8}s . As the excitation energy was increased the lifetime of the TS became $\sim 10^{-8}\text{s}$ over a substantial spectral range (1.70-1.82eV) and then, for the most energetic TS (excitation from 1.82-2.30eV) the TS lifetimes became $<10^{-8}\text{s}$. Excitation at low energies was assumed to form the TS with insufficient internal energy to access configurations in which surface hopping to the product ground state could occur.

6.2 Future directions

Future experiments are planned to explore the relative reactivity following excitation to different initial configurations of the transition state, i.e. $\text{Na}\cdots\text{XR} + hc/\lambda_1 \rightarrow [\text{Na}^*\cdots\text{XR}]^\ddagger \rightarrow \text{NaX} + \text{R}$. The detection and characterization of the products, the alkali-halide molecule, NaX, and/or the radical, CH_3 or H, by resonant-enhanced multiphoton ionization (REMPI) is being explored. The intention is to determine the yield and energy distribution in the products as a function of the excitation

wavelength, λ_1 , and hence as a function of the initial Transition State (TS) geometry. The CH_3 radicals may be probed using (3+1) REMPI[117] and the translational distribution of recoiling H atoms can be measured using Rydberg-atom time-of-flight spectroscopy[118]. This has not, however, yet been achieved for the systems described here.

Recently, a REMPI scheme for measuring the internal-energy of NaF has been proposed by this laboratory[119]. This approach was based on high level *ab initio* computations of the (ionic) ground state and the (covalent) electronically-excited states of NaF. An avoided crossing between the two states gave rise to a shallow potential-energy well in the excited state which can act as the resonant state for REMPI. This excited state would be accessed from different vibrational levels of the ground state by a photon with wavelength, $\lambda=210\text{-}340\text{nm}$. A second photon of the same wavelength would then be sufficient to ionize the NaF molecule. The potential-energy curves and the REMPI scheme are illustrated in fig. 6.1. The lifetime of the excited-state has been explored by quantum scattering of a wave-packet, and found to be $\sim 1\text{ps}$. Experience with molecules having comparable resonant-state lifetimes, such as CH_3I [120] indicates that this lifetime is sufficient for an intermediate state in a multiphoton ionization process. However, the yield of $\text{Na}\cdots\text{XR}$ complexes obtained in the crossed-molecular beam work described in this thesis has been found to be too low to detect the products NaX obtained from the photoinduced reaction.

It is believed that in the near future a larger yield of complexes can be obtained with Li metal atoms. The complex $\text{Li}\cdots\text{FH}$ has been

Figure 6.1: MRDCI calculation of the potential energy curves for the ground state, $1^1\Sigma^+$, and excited states, $n^1\Sigma^+$ and $n^1\Pi$, of NaF[119].

determined by means of high level *ab initio* calculation[84,121,122] and experiment[59] to be more strongly bound than Na·FH (by ~4×). In particular, the exchange reaction $\text{Li} + (\text{HF})_2 \rightarrow \text{Li} \cdots \text{FH} + \text{HF}$ that forms the complexes to be studied will be exoergic ($\Delta H^\circ \approx -4 \text{ kcal/mol}$ where $D^\circ(\text{Li} \cdots \text{FH}) = \sim 7 \text{ kcal/mol}$ [59,84,121,122] and $D^\circ(\text{FH} \cdots \text{FH}) = 2.97 \text{ kcal/mol}$ [111]). Consequently, a much higher concentration of the complex could be obtained which may enable an analysis of the internal-state distribution in the products.

The system Li·FH has the further advantage that it is more appropriate for accurate *ab initio* calculations due to the smaller atomic number of the alkali atom. Thus, experiment and theory can be united to give a detailed description of the dynamics of the photoinduced reaction. Quantum scattering of a wavepacket has already been studied, elsewhere, on the ground-state PES [123-126] but, as yet, the dynamics of the excited-state reaction have not been explored. A recent high level *ab initio* calculation has been made of the complete PES for the ground and first-excited state of LiFH by Paniagua and co-workers[84].

As a result of the much lower vapor pressure of Li, using an oven as the source of the alkali-atom beam will not be feasible. However, laser ablation of a metal rod can be used to generate a molecular beam of Li in an inert-carrier gas which can be directed into the expansion region of a supersonic-jet containing the halide molecules. The low collision energy which can be obtained in the absence of the oven source and using argon as the carrier gas ($< 0.02 \text{ eV}$) will be suitable for formation of Li·FH complexes. Experiments by Y. T. Lee and co-workers[60] indicated that a collision energy of at least $\sim 0.10 \text{ eV}$ is needed for reaction to occur on the

ground-state PES (i.e. in the absence of photoexcitation), hence the $\text{Li}\cdots\text{FH}$ will not be removed by thermal reaction prior to photoexcitation.

The ground-state reaction $\text{Na}+\text{FH}\rightarrow\text{NaF}+\text{H}$ is endoergic by $\sim 1.3\text{eV}$. Photodepletion of $\text{Na}\cdots\text{FH}$ complexes could be achieved by excitation of the H-F stretching mode within the complex in the electronic-ground state. If one were to excite the second overtone vibration (i.e. $v=3\leftarrow 0$), one will have supplied the complex with enough energy to surmount the energy barrier to give $\text{NaF}+\text{H}$. State-selective preparation of HF has been achieved using the infrared-radiation pumping technique[57,58]. The wavelength used to excite the vibrational mode would, of course, need to be adjusted somewhat when HF is bound in the complex. With HF vibrationally excited in $\text{Li}\cdots(\text{FH})^\dagger$ depletion of the complex would occur over a range of excitation wavelengths which correspond to different TS configurations (i.e. different $\text{Li}\cdots\text{F}$ bond stretches and $\text{Li}\cdots\text{F}-\text{H}$ angles) for the vibrationally-excited complex $\text{Li}\cdots(\text{FH})^\dagger$. Thus, the important region of the ground-state PES near the saddle point could be probed. The lifetime of $\text{Li}\cdots(\text{FH})^\dagger$ with respect to dissociation to $\text{Li}+\text{FH}$ on the ground PES is estimated to be long enough to permit the present type of photoexcitation study.

The adiabatic potential-energy curves for the system NaFH illustrated in fig.4.5 (from ref. 110) indicated a second region of avoided crossing between the ground and excited-state PES at a configuration of NaFH in which both the Na-F and H-F bond lengths are extended. The region of this second avoided crossing could be reached if the HF entity were vibrationally excited within the complex prior to formation of $[\text{Na}^*\cdots\text{FH}]^\ddagger$. This approach, in which initial vibrational excitation assists

in probing regions on an excited-state PES normally inaccessible from the ground state, has been applied to bound molecules by Crim and co-workers[127]. In their experiments the influence of particular nuclear motions on dissociation has been determined.

A more detailed understanding of the Transition State Spectroscopy presented in this thesis in terms of the evolution of $[\text{Na}^+\cdots\text{FH}]^\ddagger$ on the excited-state PES and the decay of the TS into product channels on the ground-state PES can be obtained through computation by trajectory surface-hopping studies. There are two versions of this semi-classical technique. In the Blais-Truhlar method[112], surface hopping is allowed to occur if the hopping probability is 50%. This approach is not suitable for the systems described here in which the potential energy gap between the PESs is fairly large and the hopping probabilities are much smaller. More suitable is the Tully approach[113] in which the classical trajectories are allowed to oscillate on the excited-state PES for a long time, and the hopping probability accumulates every time the region of close approach is crossed. These calculations are currently in progress at the University of Minnesota and in this laboratory. In addition, quantum scattering of a wavepacket on the excited-state PES is being studied by Zeiri and co-workers in collaboration with this laboratory. The results from these calculations will be compared with the TSH study in a future publication[128].

Thesis Summary

Van der Waals complexes, $\text{Na}\cdots\text{XR}$ ($\text{X} = \text{F}, \text{Cl}, \text{Br}$, $\text{R} = \text{CH}_3$; $\text{X} = \text{F}$, $\text{R} = \text{H}$), have been formed in a crossed-molecular beam apparatus. The Na beam was generated from an oven source and the halide beam came from the pulsed expansion of this second species in an inert-carrier gas. The concentration of complexes, $\text{Na}\cdots\text{XR}$, was determined by time-of-flight mass spectrometry (TOFMS).

Monochromatic excitation of the stable complex, $\text{Na}\cdots\text{XR}$, accessed the same potential-energy surface (PES) that would be traversed by the reaction $\text{Na}^* + \text{XR} \rightarrow \text{NaX} + \text{R}$ (where Na^* is electronically-excited Na). The excited state is formed in a selected configuration intermediate between reactants and products, for the above reaction; i.e. in the Transition State (TS) region, $[\text{Na}^*\cdots\text{XR}]^\ddagger$. By varying the excitation wavelength, the reaction could be initiated in different configurations of the TS. This constitutes a Transition State Spectroscopy for these reactions.

Charge-transfer occurs across a narrow region of configurations in the evolving TS, $[\text{Na}^*\cdots\text{XR}]^\ddagger \rightarrow [\text{Na}^+\cdots\text{XR}^-]^\ddagger$, followed by dissociation of the XR^- ion when the product, NaX, is formed. The charge-transfer process is referred to as 'harpooning'; it occurs in the region that the zero-order ionic and covalent states of the complex are degenerate.

This thesis reports measurements (referred to as 'action spectra') in which the complex $\text{Na}\cdots\text{XR}$ was photodissociated and the cross-section measured as a function of the excitation wavelength. The observation and interpretation of these action spectra for $\text{Na}\cdots\text{XR}$ has been performed

for the first time in this laboratory. We know of no other studies of these reactions using this TS spectroscopy approach.

For $\text{Na}\cdots\text{XCH}_3$ ($\text{X}=\text{F}, \text{Cl}, \text{Br}$), it was found that, by increasing the excitation frequency, ν_1 , the TS could be formed with successively higher electronically-excited states of $\text{Na}^*(i)$, i.e. $i=3p\ ^2P, 4s\ ^2S, 3d\ ^2D$. This was apparent in the photodepletion spectra for the complexes in which depletion of the stable ground-state complex, $\text{Na}\cdots\text{XCH}_3$, was measured as a function of the excitation frequency, ν_1 , or wavelength, λ_1 .

The range of wavelengths used for photoexcitation corresponded to an energy interval of $\sim 1.5\text{eV}$. The action spectra consisted of as many as four broad peaks. Each peak was assigned according to the excited state of the reactant Na atoms which correlated with the $\text{Na}^*\cdots\text{XR}$ PES. The probability with which depletion occurred on each excited PES was found to increase with electronic excitation in Na, i.e. $3p\ ^2P < 4s\ ^2S < 3d\ ^2D$ and also down the series of methyl-halide molecules, i.e. $\text{Na}\cdots\text{FCH}_3 < \text{Na}\cdots\text{ClCH}_3 < \text{Na}\cdots\text{BrCH}_3$ (this sequence corresponds to decreasing X-C bond-dissociation energy) for a particular electronically-excited state. Vibrational structure within the photodepletion peaks could be identified as being due to stretching and bending modes in the TS.

This work is believed to be the first in which the complex $\text{Na}\cdots\text{FH}$ has been observed even in its ground-electronic state. Photodepletion of this new complex has been recorded in this work across a similar energy interval for the excitation photon as was used for $\text{Na}\cdots\text{XCH}_3$. Only two peaks were observed in the action spectrum for $\text{Na}\cdots\text{FH}$. These arise (in a similar scenario to that for $\text{Na}\cdots\text{FCH}_3$) from excitation to a PES which

would be traversed by Na^* in its $3p\ 2P$ state when reacting with an HF molecule. Accurate *ab initio* calculations[110] indicate the presence of non-degenerate reactive PESs, which correlate with the $\text{Na}^*(3p\ 2P) + \text{HF}$ asymptotic limit, in the wavelength region in which photodepletion was observed in the present study.

Dissociation of the TS is believed to occur via surface hopping from the electronically-excited state into one of two possible product channels on the ground-state PES. The surface hop occurs in the region in which the character of the TS changes from largely covalent to largely ionic, $[\text{Na}^+ \cdots \text{FH}]^\ddagger \rightarrow [\text{Na}^+ \cdots \text{FH}^-]^\ddagger$. This is the harpooning event, first described over half a century ago to occur in the reaction of alkali metals with halogen molecules. The alternative outcomes are chemical reaction ($\rightarrow \text{NaF} + \text{H}$) or inelastic dissociation ($\rightarrow \text{Na} + \text{FH}^+$). At present we have no means of discriminating experimentally between these final states.

An alternative approach to Transition State Spectroscopy which provided very different information was described for the reaction $\text{Na}^*(3p\ 2P) + \text{FCH}_3 \rightarrow \text{NaF} + \text{CH}_3$. The measurement of ionization directly from the excited state, $[\text{Na} \cdots \text{FCH}_3]^\ddagger$, provided an estimate of the lifetime of the TS as a function of the wavelength used to initiate reaction in $\text{Na} \cdots \text{FCH}_3$ and therefore, in principle, as a function of the starting configuration in the TS. An action spectrum was obtained by measuring the ion signal for direct ionization from the TS as a function of the excitation energy used in forming the TS. It was found that the TS formed at low excitation energies had a lifetime which exceeded 10^{-8}s but, as the energy was increased, the lifetime became progressively shorter culminating in a lifetime of less than 10^{-8}s . At low excitation energy the TS formed has a

low probability of surface hopping and exists for a long time. In this region of long TS lifetimes, the mechanism by which the complex depletes may be laser-induced stimulated emission to dissociative configurations on the ground-state PES.

References

- (1) *Laser Spectroscopy and Photochemistry on Metal Surfaces*, ed. H.-L. Dai, and W. Ho, World Scientific, Singapore(1995).
- (2) S. K. Shin, Y. Chen, S. Nickolaisen, S. W. Sharpe, R. A. Beaudet and C. Wittig, *Advances in Photochemistry* Vol. 16, p249, ed. D. Volman, G. Hammond, and D. Neckers, Wiley Interscience, New York (1991), and references therein.
- (3) I. Harrison, J. C. Polanyi, and P. A. Young, *J. Chem. Phys.*, **89**, 1498 (1988).
- (4) St. J. Dixon-Warren, I. Harrison, K. Leggett, M. S. Matyjaszczyk, J. C. Polanyi, and P. A. Young, *J. Chem. Phys.*, **93**, 3659 (1990).
- (5) E. B. D. Bourdon, C. C. Cho, P. Das, J. C. Polanyi, C. D. Stanners, and G. Q. Xu, *J. Chem. Phys.*, **95**, 1361 (1991).
- (6) C. Jouvét, and B. Soep, *Chem. Phys. Lett.*, **96**, 426 (1983).
- (7) W. H. Breckenridge, C. Jouvét, and B. Soep, *J. Chem. Phys.*, **84**, 1443 (1986).
- (8) J. P. Visticot, B. Soep, and J. Whitham, *J. Phys. Chem.*, **92**, 4574 (1988).
- (9) B. Soep, S. Abbes, A. Keller, and J. P. Visticot, *J. Chem. Phys.*, **96**, 440 (1991).
- (10) B. Soep, C. J. Whitham, A. Keller, J. P. Visticot, *Faraday Discuss. Chem. Soc.*, **91**, 191 (1991).

- (11) A. Keller, R. Lawruszczuk, B. Soep, and J. P. Visticot, *J. Chem. Phys.*, **105**, 4556 (1996).
- (12) J. C. Polanyi, *Faraday Discuss. Chem. Soc.*, **67**, 129 (1979).
- (13) M. Polanyi, *Atomic Reactions*, Williams and Norgate, London (1932).
- (14) J. L. Magee, *J. Chem. Phys.*, **8**, 687 (1940).
- (15) D. R. Herschbach, *Adv. Chem. Phys.* **10**, 319 (1966).
- (16) K. Liu, J. C. Polanyi, and S. Yang, *J. Chem. Phys.* **96**, 8628 (1992).
- (17) K. Liu, J. C. Polanyi, and S. Yang, *J. Chem. Phys.* **98**, 5431 (1993).
- (18) J. C. Polanyi, J.-X. Wang, and S. Yang, *Isr. J. Chem.* **35**, 55 (1994).
- (19) J. C. Polanyi, and J.-X. Wang, *J. Phys. Chem.* **99**, 13691 (1995).
- (20) St. J. Dixon-Warren, K. Leggett, M. S. Matyjasczyk, J. C. Polanyi, and P. A. Young, *J. Chem. Phys.*, **88**, 4092 (1988).
- (21) St. J. Dixon-Warren, E. T. Jensen, and J. C. Polanyi, *Phys. Rev. Lett.*, **67**, 2395 (1991).
- (22) St. J. Dixon-Warren, E. T. Jensen, and J. C. Polanyi, *J. Chem. Phys.*, **98**, 5938 (1993).
- (23) J. C. Polanyi, and A. H. Zewail, *Acc. Chem. Res.* **28**, 119 (1995).
- (24) P. R. Brooks, *Chem. Rev.*, **88**, 407 (1988).
- (25) A. H. Zewail, *Faraday Discuss. Chem. Soc.* **91**, 207 (1991), and references therein.

- (26) D. G. Imre, J. K. Kinsey, R. W. Field, and D. H. Katayama, *J. Phys. Chem.*, **86**, 2564 (1982).
- (27) D. G. Imre, J. K. Kinsey, A. Sinha, and J. Krenos, *J. Phys. Chem.*, **88** 3956 (1984).
- (28) A. Weaver, and D. M. Neumark, *Faraday Discuss. Chem. Soc.* **91**, 5 (1991).
- (29) D. M. Neumark, *Acc. Chem. Res.* **26**, 33 (1993).
- (30) T. S. Rose, M. J. Rosker, A. H. Zewail, *J. Chem. Phys.*, **91** 7415 (1989).
- (31) J. H. Birely and D. R. Herschbach, *J. Chem. Phys.* **44**, 1690 (1966).
- (32) J. H. Bireley, R. R. Herm, K. R. Wilson and D. R. Herschbach, *J. Chem. Phys.* **47**, 993 (1967).
- (33) R. Grice, and P. B. Empedocles, *J. Chem. Phys.*, **48**, 5352 (1968).
- (34) J. H. Birely, E. A. Entemann, R. R. Herm, and K. R. Wilson, *J. Chem. Phys.*, **51**, 5461 (1969).
- (35) G. H. Kwei, J. A. Norris and D. R. Herschbach, *J. Chem. Phys.* **52**, 1317 (1970).
- (36) D. S. Y. Hsu and D. R. Herschbach, *Faraday Discuss. Chem. Soc.* **55**, 116 (1973).
- (37) J. C. Polanyi, *Acc. Chem. Res.*, **5**, 161 (1972); *Science*, **236**, 680 (1987).

- (38) M. G. Evans, and M. Polanyi, *Trans. Faraday Soc.*, **34**, 11 (1938).
- (39) R. B. Bernstein, A. M. Rulis, *Faraday Discuss. Chem. Soc.* **55**, 293 (1973).
- (40) G. Marcelin, and P. R. Brooks, *J. Am Chem. Soc.*, **95**, 7885 (1973); **97**, (1975).
- (41) D. H. Parker, K. K. Chakravorty, and R. B. Bernstein, *J. Phys. Chem.*, **85**, 466 (1981).
- (42) M. Trenary, H. F. Schaefer III, P. Kollman, *J. Am. Chem. Soc.*, **99**, 3885 (1977); *J. Chem. Phys.* **68**, 4047 (1978).
- (43) S. A. Pace, H. F. Pang, and R. B. Bernstein, *J. Chem. Phys.* **66**, 3635 (1977).
- (44) K. T. Wu, H. F. Pang, and R. B. Bernstein, *J. Chem. Phys.* **68**, 1064 (1978).
- (45) H. F. Pang, K. T. Wu, and R. B. Bernstein, *J. Chem. Phys.* **69**, 5267 (1978).
- (46) V. Saez Rabanos, E. Verdasco, V. J. Herrero, and A. Gonzalez Ureña, *J. Chem. Phys.*, **81**, 5725 (1984).
- (47) R. M. Harris and D. R. Herschbach, *Faraday Discuss. Chem. Soc.* **55**, 121 (1973).
- (48) K. T. Wu, *J. Phys. Chem.* **83**, 1043 (1979).

- (49) P. S. Weiss, J. M. Mestdagh, H. Schmidt, M. H. Covinsky, Y. T. Lee, *J. Phys. Chem.* **95**, 3005 (1991).
- (50) J. A. Kerr, E. A. Lissi, and A. F. Trotman-Dickenson, *J. Chem. Soc.* **64**, 1673 (1964).
- (51) P. Piecuch, *J. Mol Struct.*, **436-437**, 503 (1997).
- (52) H. H. Dispert, M. W. Geis, and P. R. Brooks, *J. Chem. Phys.*, **63**, 1173 (1975).
- (53) J. G. Pruett, F. R. Garabiner, and P. R. Brooks, *J. Chem. Phys.*, **70**, 5317 (1979).
- (54) F. Heismann, and H. J. Loesch, *Chem. Phys.* , **64**, 43 (1981).
- (55) M. Hoffmeister, L. Potthast, and H. Loesch, *Chem. Phys.* , **78**, 369 (1983).
- (56) H. J. Loesch, M. Hoffmeister, and F. Stienkemeier, *J. Chem. Soc., Faraday Trans. 2*, **85**, 1052 (1989).
- (57) H. J. Loesch, E. Stenzel, and B. Wüstenbecker, *J. Chem. Phys.*, **95**, 3841 (1991).
- (58) H. J. Loesch, and F. Stienkemeier, *J. Chem. Phys.*, **98**, 9570 (1993).
- (59) H. J. Loesch, and F. Stienkemeier, *J. Chem. Phys.*, **99**, 9598 (1993).
- (60) H. Loesch, *Chem. Phys.*, **104**, 213 (1986).
- (61) C. H. Becker, P. Casavecchia, P. W. Tiedemann, J. J. Valentini, and Y. T. Lee, *J. Chem. Phys.*, **73**, 2833 (1980).

- (62) M. F. Vernon, H. Schmidt, P. S. Weiss, M. H. Covinsky, and Y. T. Lee, *J Chem. Phys.*, **84**, 5580 (1986).
- (63) J. M. Mestdagh, B. A. Balko, M. H. Covinsky, P. S. Weiss, M. F. Vernon, H. Schmidt, and Y. T. Lee, *Faraday Discuss. Chem. Soc.*, **84**, 145 (1987).
- (64) P. S. Weiss, J. M. Mestdagh, M. H. Covinsky, B. A. Balko, and Y. T. Lee, *Chem. Phys.*, **126**, 93 (1988).
- (65) J. M. Mestdagh, and Y. T. Lee, *J. Chem. Soc., Faraday Trans. 2*, **85**, 1051 (1989).
- (66) R. Düren, U. Lackschewitz, S. Milosevic, H. Panknin and N. Schirawski, *Chem. Phys. Lett.*, **143**, 45 (1988).
- (67) R. Düren, U. Lackschewitz, S. Milosevic, *Chem. Phys.*, **126**, 81 (1988).
- (68) R. Düren, U. Lackschewitz, S. Milosevic, and H. J. Waldapfel, *J. Chem. Soc., Faraday Trans. 2*, **85**, 1017 (1989).
- (69) R. Düren, *J. Chem. Soc., Faraday Trans. 2*, **85**, 1050 (1989).
- (70) B. A. Blackwell, J. C. Polanyi, and J. J. Sloan, *Chem. Phys.*, **30**, 299 (1978).
- (71) F. E. Bartoszek, B. A. Blackwell, J. C. Polanyi, and J. J. Sloan, *J. Chem. Phys.*, **74**, 3400 (1981).
- (72) P. J. Kuntz, E. M. Nemeth, and J. C. Polanyi, *J. Chem. Phys.*, **50**, 4607 (1969).

- (73) P. J. Kuntz, M. H. Mok, and J. C. Polanyi, *J. Chem. Phys.*, **50**, 4623 (1969).
- (74) M. Shapiro, and Y. Zeiri, *J. Chem. Phys.*, **70**, 5264 (1979).
- (75) M. Paniagua, J. M Garcia de la Vega, J. R. Alvarez Collado, J. C. Sanz, J. M. Avariño, and A. Laganá, *J. Mol. Struct.*, **142**, 525 (1986).
- (76) A. Sevin, P. C. Hilberty, and J.-M. Lefour, *J. Am. Chem. Soc.*, **109**, 1845 (1987).
- (77) A. Laganá, M. L. Hernandez, J. M. Avariño, L. Castro, P. Palmieri, *Chem. Phys. Lett.*, **202**, 284 (1993).
- (78) A. Laganá, J. M. Avariño, M. L. Hernandez, P. Palmieri, E. Garcia, and T. Martinez, *J. Chem. Phys.*, **106**, 10222 (1997).
- (79) D. R. Herschbach, *Faraday Discuss. Chem. Soc.*, **62**, 162 (1977).
- (80) G. G. Balint-Kurti, and R. N. Yardley, *Faraday Discuss. Chem. Soc.*, **62**, 77 (1977).
- (81) M. M. Gallo, and D. R. Yarkony, *J. Chem. Phys.*, **86**, 4990 (1987).
- (82) K. Yamashita, and K. Morokuma, *Chem. Phys. Lett.*, **169**, 263 (1990).
- (83) D. R. Yarkony, *Int. J. Quantum Chem.*, **31**, 91 (1987).
- (84) A. Aguado, C. Suárez, and M. Paniagua, *Chem. Phys.*, **201**, 107 (1995).
- (85) M. Shapiro, and Y. Zeiri, *J. Chem. Phys.*, **85**, 6449 (1986).

- (86) C. P. Schulz, R. Haugstätter, H. U. Tittes, I. V. Hertel, *Phys. Rev. Lett.* **14**, 1704 (1986); *Z. Phys. D*, **10**, 279 (1988).
- (87) S. H. P. Bly, *Ph.D. Thesis*, Dept. of Chemistry, University of Toronto (1982).
- (88) *JANAF Thermochemical Tables*, Dow Chemical, Midland, Michigan (1965).
- (89) R. J. Gordon, Y. T. Lee, and D. R. Herschbach, *J. Chem. Phys.*, **54**, 2393 (1971).
- (90) K. Bergmann, U. Hefter, and P. Hering, *Chem. Phys.*, **32**, 329 (1978).
- (91) A. Bondi, *J. Phys. Chem.*, **68**, 441 (1964).
- (92) *Atomic and Molecular Beam Methods*, ed. G. Scoles, Oxford University Press (1988).
- (93) *CRC Handbook of Chemistry and Physics*, 77th ed.; CRC Press: Boca Raton, FL, (1996).
- (94) D. C. McGilvery, and R. J. S. Morrison, Centre for High Resolution Spectroscopy and Opto-Electronic Technology, Dept. of Chemistry, Monash University, Wellington Rd., Clayton, Victoria, 3168, Australia (Distributed by Montech Pty. Ltd.).
- (95) K. Liu, *M.Sc. Thesis*, Dept. of Chemistry, University of Toronto (1992).
- (96) X. Y. Chang, R. Ehlich, A. J. Hudson, J. C. Polanyi, and J.-X. Wang, *J. Chem. Phys.*, **106**, 3988 (1997).

- (97) H. K. Shin, *Chem. Phys. Lett.* **45**, 533 (1977).
- (98) *Alkali Halide Vapors*, ed. P. Davidovits, and D. L. McFadden, Academic Press, New York (1979).
- (99) P. Piccuch, *unpublished work*.
- (100) P. Davidovits, and D. C. Brodhead, *J. Chem. Phys.* **46**, 2668 (1967).
- (101) J. A. Kerr, *Chem. Rev.*, **66**, 465 (1966).
- (102) P. Labastie, J.-M. L'Hermite, Ph. Poncharal, and M. Sence, *J. Chem. Phys.* **103**, 6362 (1995).
- (103) C. Nitsch, Chr. Hüglin, I. V. Hertel and C. P. Schulz, *J. Chem. Phys.* **101**, 6559 (1994).
- (104) D. L. Feldman, R. K. Lengel, and R. N. Zare, *Chem. Phys. Lett.* **52**, 413 (1977).
- (105) A. Herrmann, S. Leutwyler, E. Schumacher, L. Wöste, *Chem. Phys. Lett.* **52**, 418 (1977).
- (106) C. P. Schulz, J. Höhndorf, P. Brockhaus, F. Noack, I. V. Hertel, *Chem. Phys. Lett.* **239**, 18 (1995).
- (107) R. J. Buenker, and S. D. Peyerimhoff, *Theor. Chim. Acta*, **35**, 33 (1974); **39**, 217 (1975).
- (108) P. J. Bruna, and S. D. Peyerimhoff, *Adv. Chem. Phys.* , **67**, 1 (1987).

- (109) J. J. Sakurai, *Modern Quantum Mechanics*, Addison-Wesley, Redwood City (1985).
- (110) M. S. Topaler, D. G. Truhlar, X.-Y. Chang, P. Piecuch, and J. C. Polanyi, *J. Chem. Phys.*, **108**, 5349 (1998).
- (111) A. S. Pine, and B. J. Howard, *J. Chem. Phys.*, **84**, 590 (1986).
- (112) N. C. Blais, and D. G. Truhlar, *J. Chem. Phys.*, **79**, 1334 (1983).
- (113) J. C. Tully, *J. Chem. Phys.*, **93**, 1061 (1990).
- (114) M. S. Topaler, D. G. Truhlar, X.-Y. Chang, P. Piecuch, and J. C. Polanyi, *J. Chem. Phys.*, **108**, 5378 (1998).
- (115) P. Arrowsmith, S. H. P. Bly, P. E. Carters, and J. C. Polanyi, *J. Chem. Phys.*, **79**, 283 (1983).
- (116) J. Berkowitz, *Photoabsorption, Photoionization and Photoelectron Spectroscopy*, Academic Press (1979).
- (117) J. C. Polanyi, N. S.-K. Sze, and J.-X. Wang, *Am. Chem. Soc.*, (1997).
- (118) J. B. Giorgi, R. Kühnemuth, J. C. Polanyi, and J.-X. Wang, *J. Chem. Phys.*, **106**, 3129 (1997)
- (119) X.-Y. Chang, A. J. Hudson, P. Piecuch, and J. C. Polanyi, *Faraday Discuss. Chem. Soc.*, **108**, 463 (1997).
- (120) S. P. Sapers and D. J. Donaldson, *Chem. Phys. Lett.*, **173**, 257 (1990).

- (121) P. Palmieri, and A. Laganà, *J. Chem. Phys.*, **91**, 7303 (1989).
- (122) O. N. Ventura, *Mol. Phys.*, **89**, 1851 (1996).
- (123) F. Gögtas, G. G. Balint-Kurti, A. R. Offer, *J. Chem. Phys.*, **104**, 7927 (1996).
- (124) A. Aguado, M. Paniagua, M. Lara, and O. Roncero, *J. Chem. Phys.*, **106**, 1013 (1997).
- (125) W. Zhu, D. Wang, and J. Z. H. Zhang, *Theor. Chem. Acc.*, **96**, 31 (1997).
- (126) J. M. Alvariño, V. Aquilanti, S Cavalli, S Crocchianti, A Laganà, and T. Martínez, *J. Chem. Phys.*, **107**, 3339 (1997).
- (127) F. F. Crim, *Annu. Rev. Phys. Chem.*, **44**, 397 (1993).
- (128) M. S. Topaler, D. G. Truhlar, P. Piecuch, and J. C. Polanyi, in *preparation*.